

Planetesimal disk evolution driven by planetesimal-planetesimal gravitational scattering.

R. R. Rafikov

Princeton University Observatory, Princeton, NJ 08544

rrr@astro.princeton.edu

ABSTRACT

We investigate the process of an inhomogeneous planetesimal disk evolution caused by the planetesimal-planetesimal gravitational scattering. We develop a rather general approach based on the kinetic theory which self-consistently describes the evolution in time and space of both the disk surface density and its kinematic properties — dispersions of eccentricity and inclination. The gravitational scattering of planetesimals is assumed to be in the dispersion-dominated regime which considerably simplifies analytical treatment. The resultant equations are of advection-diffusion type. Distance dependent scattering coefficients entering these equations are calculated analytically under the assumption of two-body scattering in the leading order in Coulomb logarithm. They are essentially nonlocal in nature. Our approach allows one to explore the dynamics of nonuniform planetesimal disks with arbitrary mass and random velocity distributions. It can also naturally include other physical mechanisms which are important for the evolution of such disks — gas drag, migration, and so on.

Subject headings: planets and satellites: general — solar system: formation — (stars:) planetary systems

1. Introduction.

The discovery of extrasolar giant planets around nearby stars (Mayor & Queloz 1995; Marcy *et al.* 2000; Vogt *et al.* 2000; Butler *et al.* 2001) has been one of the most exciting astrophysical findings of the last decade. It has revived interest in planetary sciences and stimulated many theoretical studies. One of the areas which has received a lot of attention recently is the formation of terrestrial planets — planets like Earth or Venus.

There are many reasons for this interest. First of all, our own Solar System hosts several terrestrial planets and we must understand their formation mechanisms if we want to know the history of our planetary system. Second, despite the huge differences in their physical properties, giant planets are probably linked to the terrestrial ones through their formation mechanism, since it is widely believed that giant planets form as a result of gas accretion on a preexisting rocky core

which essentially was a massive Earth-type planet (Mizuno 1980; Stevenson 1982; Bodenheimer & Pollack 1986; see however Boss *et al.* 2002). Thus terrestrial planets are probably an important component in the evolutionary history of the giant planets. Third, a lot of effort is currently being directed toward designing and building space missions with the ultimate goal of detecting Earth-type planets around other stars. Theoretical understanding of how terrestrial planets had come into being will help us plan these missions most effectively. Finally, observations of IR emission from the dust and debris disks around nearby stars (Heinrichsen *et al.* 1999; Schneider 2001) must be interpreted in the context of the formation and evolution of such disks. They are very likely to be the outcome of the same collisional fragmentation and accumulation of massive rocky or icy bodies that form terrestrial planets.

It is widely believed that the formation of Earth-type planets proceeded via agglomeration of large number of planetesimals — asteroid or comet-like rocky or icy bodies. The theory of this process was pioneered by Safronov (1972) who was the first to point out the importance of the evolution of the dynamical properties of the planetesimal disk for the evolution of its mass distribution. The discovery of a rapid “runaway” mode of planetary accretion by Wetherill & Stewart (1989) has made this issue even more important since this mechanism relies on the redistribution of the energy of planetesimal epicyclic motion between populations with different masses (Wetherill & Stewart 1993; Kenyon & Luu 1998; Inaba *et al.* 2001). Significant progress has been made recently in understanding the dynamical evolution of *homogeneous* planetesimal disks — disks where gradients of surface density or dynamical properties (such as the dispersions of eccentricity and inclination of various planetesimal populations) are absent (Ida 1990; Stewart & Ida 2000, hereafter SI; Ohtsuki *et al.* 2002). This assumption should be very good during the initial stages of planetesimal growth when there are no massive bodies in the disk. However as coagulation proceeds and planetary embryos — precursors of terrestrial planets — emerge this assumption runs into problems. It was first demonstrated by Ida & Makino (1993) using N-body simulations and then confirmed using orbit integrations (Tanaka & Ida 1997, 1999) that under a variety of conditions massive protoplanetary embryos would tend to repel planetesimal orbits, clearing out an annular gap in the disk around them. This process introduces a new spatial dimension into the problem and makes it much harder to treat.

N-body simulations are not good tools to study the details of this process. The primary reason is that they become too time-consuming when one needs to follow the spatial and dynamical properties of a many-body system for many orbital periods. Moreover, the number of planetesimals which they can handle is not very large (less than 10^4) which precludes the consideration of realistic planetesimal disks containing huge number of bodies with masses spanning an enormous range — from one meter rocks to 100 km planetesimals. Finally, one would need to perform a large number of such simulations to explore the whole space of physical parameters relevant for protoplanetary disk evolution. Similar problems, although less severe, plague the performance of methods based on the direct integration of planetesimal orbits using some simplifying assumptions (Tanaka & Ida 1997, 1999).

The large number of bodies under consideration poses no problem for the methods of kinetic theory. In this approach the disk is split into a set of planetesimal populations and each of them is characterized by three functions of position: the surface density and dispersions of eccentricity and inclination. Application of this method for studying planetesimal coagulation and the evolution of dynamical properties in homogeneous disks has proven to be very useful (Wetherill & Stewart 1989; Kenyon & Luu 1998). It can also be easily extended to the case of inhomogeneous disks by allowing the planetesimal properties to vary within the disk. This approach allows one to study many physical mechanisms important for the coagulation process — gravitational interactions between planetesimals increasing their random motion, gas drag and inelastic collisions damping them, migration, fragmentation, etc. The use of this statistical approach for exploring inhomogeneous disks was first undertaken by Petit & Hénon (1987a, 1987b, 1988) in their studies of planetary rings. Physical conditions in planetary rings (high optical depth and frequent inelastic collisions) are very different from those which are thought to exist in protoplanetary disks. Thus we cannot directly apply the methods of Petit & Hénon to study planetesimal disks, but the spirit in which their investigation was carried out can be largely preserved.

In this paper we consider the evolution of *inhomogeneous* planetesimal disks caused by interactions between planetesimals using conventional methods of statistical mechanics (Lifshitz & Pitaevskii 1981). The effects of the planetary embryos on the disk evolution will be investigated later in Rafikov (2002; hereafter Paper II). We will concentrate on one of the most important processes going on in these disks — mutual gravitational scattering of planetesimals — although our rather general approach allows one to treat other relevant phenomena as well. This process can proceed in two distinct regimes depending on the amplitude of planetesimal random motion. The gravitational attraction between two planetesimals with masses m_1 and m_2 becomes stronger than the tidal field of the central star of mass M_c when their mutual separation is less than their Hill (or tidal, or Roche) radius r_H , defined as

$$r_H = a_0 \left(\frac{m_1 + m_2}{M_c} \right)^{1/3}, \quad (1)$$

where a_0 is the distance from the central star. When the random velocities of the epicyclic motion of interacting planetesimals are smaller than $\sim \Omega r_H$ ($\Omega = \sqrt{GM_c/a_0^3}$ is the disk orbital frequency at a_0) their relative approach velocities are small and close interactions can lead to a considerable change of the orbital elements of planetesimals. This velocity regime is called shear-dominated (or “cold”). It should be contrasted with the other extreme — the so called dispersion-dominated (“hot”) regime which occurs when planetesimal velocity dispersions are bigger than $\sim \Omega r_H$. In this latter case scattering is typically weak which often allows analytical treatment of this velocity regime.

The development of planetesimal disk inhomogeneities driven by a protoplanetary embryo was explored in Rafikov (2001; hereafter Paper I) assuming that shear-dominated scattering of planetesimals prevails. It was also assumed in this study that the dynamical properties of planetesimals do not evolve as a result of scattering and that the disk always stays dynamically cold. Planetesimal-

planetesimal interactions play the role of effective viscosity in the disk, and tend to homogenize it and close up any gap. Nevertheless, this study demonstrated that gap formation is the natural outcome of the embryo-planetesimal interaction when the embryo is massive enough. These interactions were local in character because in the shear-dominated case only planetesimals on orbits separated by no more than several r_H (corresponding to their encounter) were able to approach each other closely.

It is however more likely that the planetesimal-planetesimal gravitational scattering in realistic protoplanetary disks proceeds in the dispersion-dominated (rather than shear-dominated) regime, at least on the late stages of disk evolution (see §3). In this case the evolution of planetesimal random motion can strongly affect the growth rate of protoplanetary embryos. It is also tightly coupled to the evolution of spatial distribution of planetesimals because any change of the energy of epicyclic motion comes at the expense of the orbital energy of planetesimals.

The treatment of the dispersion-dominated case is complicated by the fact that planetesimals in this regime can explore different regions of the disk in the course of their epicyclic motion. This makes disk evolution a nonlocal process. On the other hand, as we have said before there are natural simplifications which are valid in the dispersion-dominated regime. These include the two-body scattering approximation (relative velocities are high), Fokker-Planck type expansions (scattering is weak), and so on.

Our paper is organized as follows. In §2 we derive general equations for the planetesimal surface density and velocity dispersion evolution. We do this using the Hill approximation which is briefly described in §2.1. A Fokker-Planck expansion of the evolution equations, valid in the dispersion-dominated regime, is performed in §3. In §4 we derive analytical expressions for the scattering coefficients used in these equations. We conclude with a brief summary of our results in §5. Some technical details of the calculations and derivations can be found in appendices.

2. Equations of evolution of planetesimal disk properties.

Throughout the paper we will assume disk to be axisymmetric. All the following calculations also assume a disk with a Keplerian rotation law, although they could be easily extended to the case of an arbitrary rotation law.

As we have mentioned before planetesimal disks must have contained a huge number of constituent bodies with a wide range of masses. Thus we use a kinetic approach and characterize the state of the disk at any moment of time t by a set of three averaged parameters for each mass population: the surface number density of planetesimals and their velocity dispersions in the horizontal and vertical directions at every point $\mathbf{r}$ in the disk.

When the orbit's eccentricity and inclination are small, it is convenient to work in the guiding center approximation, when the elliptic motion is represented as an epicyclic motion of the body

on a small ellipse in horizontal plane with the radial dimension ae (and oscillatory motion in the vertical direction with the amplitude ai); the center of this ellipse performs circular motion with a semimajor axis a . In the case of Keplerian rotation law frequencies of all these motions are the same and equal to the angular frequency of Keplerian motion. In this approximation the roles of longitude of ascending node and argument of pericentre are played by two constant phase angles ω and τ which characterize the position of body on its epicycle at a given moment of time. Random motion is defined as the motion relative to the circular orbit of the guiding center, i.e. is represented by the epicyclic motion. Eccentricity and inclination thus measure the magnitude of the random motion and we will take them to represent planetesimal's kinematic properties.

The spatial distribution of planetesimals can be characterized by two kinds of surface density. One is the *surface density of guiding centers* of planetesimals $N(a)$ and is defined such that $N(a)2\pi a da$ is the number of planetesimals with guiding centers between a and $a + da$. It is a function of planetesimal semimajor axes only.

Another parametrization is the *instantaneous surface density* $N^{inst}(r)$, defined such that $N^{inst}(r)2\pi r dr$ is the number of planetesimals within the disk surface element $2\pi r dr$ (we will denote instantaneous distance from the central star as r to distinguish it from a). The relation between these two surface densities can only be computed if the distribution of planetesimal eccentricities is known (we calculate an appropriate conversion between N and N^{inst} in Appendix A). These surface densities coincide only when planetesimals are on circular orbits (which was the case in Paper I). We will mostly work with the surface density of the guiding centers and not the instantaneous surface density.

2.1. Hill equations.

In the epicyclic approximation, which is valid when $e \ll 1$ and $i \ll 1$, it is convenient to introduce the Cartesian coordinate system x, y, z with axes in the r, φ, z directions of the corresponding cylindrical system and origin at some reference distance a_0 from the central star. This system rotates around the central star with angular velocity $\Omega_0 = \Omega(a_0)$. In these coordinates the unperturbed motion of the body is given by (Goldreich & Lynden-Bell 1965; Hénon & Petit 1986)

$$\begin{aligned} x &= ha_0 - ea_0 \cos(\Omega_0 t + \tau) + O(e^2, i^2), \\ y &= -\frac{3}{2}ha_0(\Omega_0 t - \lambda) + 2ea_0 \sin(\Omega_0 t + \tau) + O(e^2, i^2), \\ z &= ia_0 \sin(\Omega_0 t + \omega) + O(e^2, i^2), \end{aligned} \tag{2}$$

where τ and ω are the constant phase angles we have mentioned before and λ is the origin of time (it is usually unimportant and can be set equal to 0). Dimensionless relative semimajor axis a of the body (the location of its guiding center) h is defined as

$$h = \frac{a - a_0}{a_0} \tag{3}$$

Instead of the pairs e, τ and i, ω it is sometimes convenient to use the eccentricity and inclination *vectors* $\mathbf{e}$ and $\mathbf{i}$:

$$\begin{aligned}\mathbf{e} &= (e_x, e_y) = (e \cos \tau, e \sin \tau), \\ \mathbf{i} &= (i_x, i_y) = (i \cos \omega, i \sin \omega).\end{aligned}\tag{4}$$

We can also use other natural simplifications. One of them is motivated by the assumption that the masses of planetesimals m_j are much smaller than the mass of the central star M_c — a condition which is always fulfilled in planetesimal disks. In this case it was demonstrated by Hénon & Petit (1986) that the equations describing the 3-body interaction (star and planetesimals 1 and 2) can be significantly simplified. In particular they have demonstrated that relative motion of interacting bodies ($\mathbf{r} = \mathbf{r}_1 - \mathbf{r}_2$) and the motion of their barycenter [$\mathbf{r}_b = (m_1 \mathbf{r}_1 + m_2 \mathbf{r}_2)/(m_1 + m_2)$] separate from each other. In the course of 3-body interactions the barycenter motion is not affected at all and remains in unperturbed epicyclic motion at all times.

We define the relative Hill coordinates of the two interacting bodies $(\tilde{x}, \tilde{y}, \tilde{z})$ with masses $m_1 \ll M_c$ and $m_2 \ll M_c$ as

$$\tilde{x} = (x_1 - x_2)/r_H, \quad \tilde{y} = (y_1 - y_2)/r_H, \quad \tilde{z} = (z_1 - z_2)/r_H.\tag{5}$$

Here r_H is the Hill radius defined by equation (1). Instead of m we will often use $\mu = m/M_c$ [with this notation the Hill radius can be written as $r_H = a_0(\mu_1 + \mu_2)^{1/3}$]. Note that the Hill radius is only meaningful for a *pair* of bodies and its definition is similar to the definition of tidal radius. We will also define *reduced* eccentricity and inclination vectors $\tilde{\mathbf{e}} = a_0(\mathbf{e}_1 - \mathbf{e}_2)/r_H$, $\tilde{\mathbf{i}} = a_0(\mathbf{i}_1 - \mathbf{i}_2)/r_H$ and the reduced impact parameter $\tilde{h} = a_0(h_1 - h_2)/r_H$.

With these definitions the equations of relative motion can be written as (Hénon & Petit 1986; Hasegawa & Nakazawa 1990)

$$\frac{d^2 \tilde{x}}{dT^2} - 2 \frac{d\tilde{y}}{dT} - 3\tilde{x} = -\frac{\partial \phi_p}{\partial \tilde{x}},\tag{6}$$

$$\frac{d^2 \tilde{y}}{dT^2} + 2 \frac{d\tilde{x}}{dT} = -\frac{\partial \phi_p}{\partial \tilde{y}},\tag{7}$$

$$\frac{d^2 \tilde{z}}{dT^2} + \tilde{z} = -\frac{\partial \phi_p}{\partial \tilde{z}},\tag{8}$$

where ϕ_p is a (dimensionless) potential of interaction between the two bodies and $T = \Omega_0 t$. For the Newtonian gravitational potential between planetesimals

$$\phi_p(\mathbf{r}) = -(\tilde{x}^2 + \tilde{y}^2 + \tilde{z}^2)^{-1/2}.\tag{9}$$

An important feature of these equations is that they do not contain any parameters of the interacting bodies (such as their masses, etc.). All the physically important information is embedded into the definitions of Hill radius and relative coordinates. Also, the outcome of the interaction between

the two bodies depends only on their *relative* orbital parameters. This universality is a very useful property which we will widely exploit later on. Equations (6)-(8) possess an integral of motion — the Jacobi constant

$$J = \left(\frac{d\tilde{x}}{dT} \right)^2 + \left(\frac{d\tilde{y}}{dT} \right)^2 + \left(\frac{d\tilde{z}}{dT} \right)^2 - 3\tilde{x}^2 + \tilde{z}^2 + 2\phi_p. \quad (10)$$

Conservation of this quantity poses an important constraint on the changes of the orbital elements of interacting bodies in the course of their gravitational interaction.

One can easily see that when the interaction is absent equations (2) represent a solution of the system (6)-(8) if we replace all quantities in (2) by their relative (and scaled by r_H) analogs. But in the presence of the interaction, orbital parameters are no longer constants and will evolve with time. However, we can still represent epicyclic motion by equations (2) keeping in mind that $\tilde{\mathbf{e}}, \tilde{\mathbf{i}}, \tilde{h}$, etc. are now *instantaneous* osculating values of the orbital elements. Their evolution is governed by the following set of equations (Goldreich & Tremaine 1980; Hasegawa & Nakazawa 1990):

$$\begin{aligned} \frac{d\tilde{h}}{dT} &= -2\frac{\partial\phi_p}{\partial\tilde{y}}, & \frac{d\lambda}{dT} &= \frac{4}{3\tilde{h}}\frac{\partial\phi_p}{\partial\tilde{x}} - \frac{2}{\tilde{h}}(T - \lambda)\frac{\partial\phi_p}{\partial\tilde{y}}, \\ \frac{d\tilde{e}_x}{dT} &= -\sin T\frac{\partial\phi_p}{\partial\tilde{x}} - 2\cos T\frac{\partial\phi_p}{\partial\tilde{y}}, & \frac{d\tilde{e}_y}{dT} &= -\cos T\frac{\partial\phi_p}{\partial\tilde{x}} + 2\sin T\frac{\partial\phi_p}{\partial\tilde{y}}, \\ \frac{d\tilde{i}_x}{dT} &= -\cos T\frac{\partial\phi_p}{\partial\tilde{z}}, & \frac{d\tilde{i}_y}{dT} &= \sin T\frac{\partial\phi_p}{\partial\tilde{z}}. \end{aligned} \quad (11)$$

The relative velocity $\dot{\mathbf{r}}$ of planetesimal motion in Hill coordinates (dimensional) is given by [see (2)]

$$\dot{x} = \tilde{e}r_H \sin(T + \tau), \quad \dot{y} = -\frac{3}{2}\tilde{h}r_H + 2\tilde{e}r_H \cos(T + \tau), \quad \dot{z} = \tilde{i}r_H \cos(T + \omega). \quad (12)$$

In many applications it is better to use the velocity defined *relative to the local velocity of the circular motion*. Clearly this velocity $\mathbf{v}$ is related to $\dot{\mathbf{r}}$ as

$$\mathbf{v} = \dot{\mathbf{r}} + \frac{3}{2}x\mathbf{n}_y, \quad (13)$$

where $\mathbf{n}_y$ is a unit vector in the y -direction. This definition implies that

$$v_x = \dot{x} = \tilde{e}r_H \sin(T + \tau), \quad v_y = \dot{y} + \frac{3}{2}x = \frac{\tilde{e}r_H}{2} \cos(T + \tau), \quad v_z = \dot{z} = \tilde{i}r_H \cos(T + \omega). \quad (14)$$

With the use of equations (2) and (14) one can express e, i and h as functions of $\mathbf{r}$ and $\mathbf{v}$ which will later be used in §4. Using (2) and (12) we can rewrite the expression (10) for the Jacobi constant in the following form:

$$J = \tilde{e}^2 + \tilde{i}^2 - \frac{3}{4}\tilde{h}^2 + 2\phi_p \quad (15)$$

(our definition of the Jacobi constant may differ from others by a constant additive and/or multiplicative factors, cf. Hasegawa & Nakazawa 1990).

2.2. Derivation of the master equation for the distribution function.

Based on the previous discussion it is natural to describe the position and velocity state of a planetesimal with mass m_j by its guiding center radius h_j (relative to the reference radius a_0), vector eccentricity $\mathbf{e}_j$ and inclination $\mathbf{i}_j$ ¹ (note that this is not a canonical set of variables!). The five-dimensional vector $\mathbf{\Gamma}_j = (h_j, \mathbf{e}_j, \mathbf{i}_j)$ will be used to concisely denote this state and $d\mathbf{\Gamma}_j = dh_j d\mathbf{e}_j d\mathbf{i}_j$ will denote a phase space volume element around $\mathbf{\Gamma}_j$.

We set $N(m, h, t)dm$ to be the dimensionless surface number density of planetesimal guiding centers in the mass interval $(m, m + dm)$. It differs from its dimensional analog $N(a)$ by a factor of a_0^2 . We introduce the notion of a planetesimal distribution function (PIDF) $f(m, \mathbf{\Gamma}, t)$ such that $2\pi f(m, \mathbf{\Gamma}, t)d\mathbf{\Gamma}dm$ is the number of planetesimals in phase space volume $d\mathbf{\Gamma}$ and in mass interval $(m, m + dm)$. Clearly,

$$\int \int f(m, \mathbf{\Gamma}, t) d\mathbf{\Gamma} dm = N(m, h, t), \quad (16)$$

where the integration is performed over the 4-dimensional space of vector eccentricity. For the purpose of brevity we will further use $f_j(\mathbf{\Gamma}, t)$ and $N_j(h, t)$ instead of $f(m_j, \mathbf{\Gamma}_j, t)$ and $N(m_j, h, t)$.

In what follows we will consider the distribution function of planetesimal vector eccentricities and inclinations to be Gaussian (or Rayleigh, or Schwarzschild). In other words, we assume that $f_j(\mathbf{\Gamma})d\mathbf{\Gamma} = N(h)\psi_j(\mathbf{e}, \mathbf{i}, h)d\mathbf{\Gamma}$, where

$$\psi_j(\mathbf{e}, \mathbf{i}, h) = \frac{1}{4\pi^2 \sigma_{ej}^2(h) \sigma_{ij}^2(h)} \exp \left[-\frac{\mathbf{e}^2}{2\sigma_{ej}^2(h)} - \frac{\mathbf{i}^2}{2\sigma_{ij}^2(h)} \right]. \quad (17)$$

Here σ_{ej}, σ_{ij} are the r.m.s. eccentricities and inclinations of planetesimals of mass m_j at point h , given by

$$\sigma_{ej} = \langle \mathbf{e}_j^2 \rangle^{1/2} / \sqrt{2}, \quad \sigma_{ij} = \langle \mathbf{i}_j^2 \rangle^{1/2} / \sqrt{2}. \quad (18)$$

Greenzweig & Lissauer (1992) and Ida & Makino (1992) have demonstrated that this assumption of Gaussian distribution is very good in their N-body simulations for large values of planetesimal eccentricities and inclinations [see also Ohtsuki (1999) for the discussion of the validity of distribution (17) in circumplanetary disks].

Consider now planetesimals with masses m_1 (type 1) and m_2 (type 2) (we usually denote the particle under consideration by subscript 1 and scattered particles by subscript 2). Scattering can bring planetesimals of type 1 into the phase volume element $d\mathbf{\Gamma}$ and can also scatter particles out of it. The number of encounters in time dt between particles of type 1 *initially* in the phase volume element $d\mathbf{\Gamma}_a$ and those of type 2 initially in $d\mathbf{\Gamma}_b$ is

$$dN_{coll} = \nu(\mathbf{\Gamma}_a, \mathbf{\Gamma}_b) f_1(\mathbf{\Gamma}_a) f_2(\mathbf{\Gamma}_b) d\mathbf{\Gamma}_a d\mathbf{\Gamma}_b dt, \quad (19)$$

¹These variables are dimensionless but not scaled by r_H as in §2.1.

where $\nu(\mathbf{\Gamma}_a, \mathbf{\Gamma}_b)$ is the rate of encounters.

Let us introduce the differential probability $P(\mathbf{\Gamma}_a, \mathbf{\Gamma}_b, \Delta\mathbf{\Gamma}_a)$ such that $P(\mathbf{\Gamma}_a, \mathbf{\Gamma}_b, \Delta\mathbf{\Gamma}_a)d(\Delta\mathbf{\Gamma}_a)$ is the probability that particle 1 changes its state from $\mathbf{\Gamma}_a$ to the region $d\Delta\mathbf{\Gamma}_a$ around $\mathbf{\Gamma}_a + \Delta\mathbf{\Gamma}_a$ in an encounter with particle of type 2, if before the collision they were in states $\mathbf{\Gamma}_a, \mathbf{\Gamma}_b$ respectively. We can then write down the evolution equation for PPDF of particles of type 1:

$$\begin{aligned} \frac{\partial f_1(\mathbf{\Gamma})}{\partial t} + \dot{\mathbf{r}} \frac{\partial f_1(\mathbf{\Gamma})}{\partial \mathbf{\Gamma}} &= \int_0^\infty dm_2 \int d\mathbf{\Gamma}_a d\mathbf{\Gamma}_b \\ &\times \left[\nu(\mathbf{\Gamma}_a, \mathbf{\Gamma}_b) f_1(\mathbf{\Gamma}_a) f_2(\mathbf{\Gamma}_b) P(\mathbf{\Gamma}_a, \mathbf{\Gamma}_b, \mathbf{\Gamma} - \mathbf{\Gamma}_a) - \nu(\mathbf{\Gamma}, \mathbf{\Gamma}_b) f_1(\mathbf{\Gamma}) f_2(\mathbf{\Gamma}_b) P(\mathbf{\Gamma}, \mathbf{\Gamma}_b, \mathbf{\Gamma}_a - \mathbf{\Gamma}) \right]. \end{aligned} \quad (20)$$

Here the second term in the l.h.s. represents evolution caused by reasons other than mutual gravitational scattering (e.g. gas drag, migration, non-Keplerian gravitational forces, etc.). The r.h.s. of (20) is the collision integral for scattering between planetesimals; its first term represents particles entering $d\mathbf{\Gamma}$, while the second one represents planetesimals leaving $d\mathbf{\Gamma}$. This is the most general form of evolution equation, which will be further simplified using additional assumptions.

2.3. Local approximation.

When the masses of the interacting bodies are much smaller than M_c their mutual Hill radius is much smaller than the distance to the central star a_0 . We will also assume that the sizes of epicycles of individual bodies are small compared to a_0 . As a result, we can safely use the Hill formalism (§2.1) to describe planetesimal scattering.

The local character of the interaction means that the encounter rate $\nu(\mathbf{\Gamma}_a, \mathbf{\Gamma}_b)$ is determined by the local shear and can be expressed as

$$\nu(\mathbf{\Gamma}_a, \mathbf{\Gamma}_b) = |\Omega(h_a) - \Omega(h_b)| = 2|A||h_a - h_b|, \quad (21)$$

where $A = (r/2)(d\Omega/dr)$ is a function characterizing shear in the disk (Oort's A constant), $A = -(3/4)\Omega_0$ for a Keplerian rotation law. Using this expansion we can rewrite equation (20) as

$$\begin{aligned} \frac{\partial f_1(\mathbf{\Gamma})}{\partial t} + \dot{\mathbf{r}} \frac{\partial f_1(\mathbf{\Gamma})}{\partial \mathbf{\Gamma}} &= 2|A| \int_0^\infty dm_2 \int d\mathbf{\Gamma}_a d\mathbf{\Gamma}_b \\ &\times \left[|h_a - h_b| f_1(\mathbf{\Gamma}_a) f_2(\mathbf{\Gamma}_b) P(\mathbf{\Gamma}_a, \mathbf{\Gamma}_b, \mathbf{\Gamma} - \mathbf{\Gamma}_a) - |h - h_b| f_1(\mathbf{\Gamma}) f_2(\mathbf{\Gamma}_b) P(\mathbf{\Gamma}, \mathbf{\Gamma}_b, \mathbf{\Gamma}_a - \mathbf{\Gamma}) \right]. \end{aligned} \quad (22)$$

We can easily integrate the second term over $d\mathbf{\Gamma}_a$ because it is clear that

$$\int d\mathbf{\Gamma}_a P(\mathbf{\Gamma}, \mathbf{\Gamma}_b, \mathbf{\Gamma}_a - \mathbf{\Gamma}) = 1, \quad (23)$$

since this integral represents the probability of scattering planetesimal 1 (initially at $\mathbf{\Gamma}$) *anywhere* in the course of interaction with planetesimal 2 (initially at $\mathbf{\Gamma}_b$). Then we can also integrate $f_2(\mathbf{\Gamma}_2)$ over $d\mathbf{e}_2 d\mathbf{i}_2$ to find

$$\int d\mathbf{\Gamma}_a d\mathbf{\Gamma}_b |h - h_b| f_1(\mathbf{\Gamma}) f_2(\mathbf{\Gamma}_b) P(\mathbf{\Gamma}, \mathbf{\Gamma}_b, \mathbf{\Gamma}_a - \mathbf{\Gamma}) = \int_{-\infty}^{\infty} dh_b |h - h_b| f_1(\mathbf{\Gamma}) N_2(h_b). \quad (24)$$

To deal with the first term in the r.h.s. of (22) we first recall that in the Hill approximation all scattering properties depend only on *relative* coordinates and velocities of planetesimals. This means that

$$P(\mathbf{\Gamma}_a, \mathbf{\Gamma}_b, \Delta\mathbf{\Gamma}_a) = P(\mathbf{\Gamma}_a - \mathbf{\Gamma}_b, \Delta\mathbf{\Gamma}_a), \quad (25)$$

Now, let us introduce the new relative variable

$$\tilde{\mathbf{\Gamma}}_r = (\tilde{h}, \tilde{\mathbf{e}}_r, \tilde{\mathbf{i}}_r) = \frac{\mathbf{\Gamma}_a - \mathbf{\Gamma}_b}{(\mu_1 + \mu_2)^{1/3}} = \frac{1}{(\mu_1 + \mu_2)^{1/3}} (h_a - h_b, \mathbf{e}_a - \mathbf{e}_b, \mathbf{i}_a - \mathbf{i}_b), \quad (26)$$

where $\mathbf{\Gamma}_a$ and $\mathbf{\Gamma}_b$ characterize the orbital elements of particles 1 and 2 correspondingly. The change of $\tilde{\mathbf{\Gamma}}_r$ in the course of an encounter is $\Delta\tilde{\mathbf{\Gamma}}_r$, and depends only on $\mathbf{\Gamma}_a$ and $\mathbf{\Gamma}_b$ through the combination $\tilde{\mathbf{\Gamma}}_r$. Then, bearing in mind conservation of the center of mass coordinate $m_1\mathbf{\Gamma}_a + m_2\mathbf{\Gamma}_b$, we obtain that $\mathbf{\Gamma}_a, \mathbf{\Gamma}_b$ can be expressed in terms of $\mathbf{\Gamma}$ [in the first term of (22) this is the value of $\mathbf{\Gamma}_a$ after the encounter], $\tilde{\mathbf{\Gamma}}_r$, and $\Delta\tilde{\mathbf{\Gamma}}_r$ as

$$\mathbf{\Gamma}_a = \mathbf{\Gamma} - \frac{\mu_2}{\mu_1 + \mu_2} (\mu_1 + \mu_2)^{1/3} \Delta\tilde{\mathbf{\Gamma}}_r, \quad \mathbf{\Gamma}_b = \mathbf{\Gamma} - (\mu_1 + \mu_2)^{1/3} \left(\frac{\mu_2}{\mu_1 + \mu_2} \Delta\tilde{\mathbf{\Gamma}}_r + \tilde{\mathbf{\Gamma}}_r \right). \quad (27)$$

Instead of the probability $P(\mathbf{\Gamma}_a, \mathbf{\Gamma}_b, \mathbf{\Gamma} - \mathbf{\Gamma}_a) d(\Delta\mathbf{\Gamma}_a)$ we will work with the probability of scattering in relative coordinates $\tilde{P}_r(\tilde{\mathbf{\Gamma}}_r, \Delta\tilde{\mathbf{\Gamma}}_r) d(\Delta\tilde{\mathbf{\Gamma}}_r)$. From the equation (27) one can easily find that

$$\tilde{P}_r(\tilde{\mathbf{\Gamma}}_r, \Delta\tilde{\mathbf{\Gamma}}_r) = \frac{\mu_2^5}{(\mu_1 + \mu_2)^{10/3}} P(\mathbf{\Gamma}_a - \mathbf{\Gamma}_b, \Delta\mathbf{\Gamma}_a). \quad (28)$$

This new probability function is independent of the masses of the interacting particles. It also follows from the properties of the Hill equations that

$$\tilde{P}_r(\tilde{\mathbf{\Gamma}}_r, \Delta\tilde{\mathbf{\Gamma}}_r) = \delta \left[\Delta\tilde{\mathbf{\Gamma}}_r - \Delta\tilde{\mathbf{\Gamma}}_{sc}(\tilde{\mathbf{\Gamma}}_r) \right], \quad (29)$$

i.e. the initial state of the scattering planetesimals unambiguously determines their final state through the scattering function $\Delta\tilde{\mathbf{\Gamma}}_{sc}(\tilde{\mathbf{\Gamma}}_r)$.

Using these definitions we can finally put the PLDF evolution equation (22) in the following form:

$$\begin{aligned} \frac{\partial f_1(\mathbf{\Gamma})}{\partial t} + \frac{d\mathbf{\Gamma}}{dt} \frac{\partial f_1(\mathbf{\Gamma})}{\partial \mathbf{\Gamma}} &= -2|A| \int_0^\infty dm_2 (\mu_1 + \mu_2)^{2/3} \int_{-\infty}^\infty d\tilde{h} |\tilde{h}| \\ &\times \left[f_1(\mathbf{\Gamma}) N_2(h - (\mu_1 + \mu_2)^{1/3} \tilde{h}) \right. \\ &\left. - (\mu_1 + \mu_2)^{4/3} \int d\tilde{\mathbf{e}}_r d\tilde{\mathbf{i}}_r d(\Delta\tilde{\mathbf{\Gamma}}_r) f_1(\mathbf{\Gamma}_a) f_2(\mathbf{\Gamma}_b) P_r(\tilde{\mathbf{\Gamma}}_r, \Delta\tilde{\mathbf{\Gamma}}_r) \right], \end{aligned} \quad (30)$$

with $\mathbf{\Gamma}, \mathbf{\Gamma}_a, \mathbf{\Gamma}_b, \tilde{\mathbf{\Gamma}}_r$ and $\Delta\tilde{\mathbf{\Gamma}}_r$ related by equations (26) and (27).

We will use this general equation for the evolution of the PIDF to obtain formulae describing the behavior of the spatial and kinematic properties of planetesimal disk.

2.4. Evolution of surface density.

In this paper we will be mostly interested in the evolution of the planetesimal disk due to the gravitational scattering between planetesimals. Thus, we set the second term in the l.h.s. of equation (30) to 0 (it can be easily reinstated when the need arises). At this stage we also replace subscripts “a” and “b” with “1” and “2” since now $\mathbf{\Gamma}_a$ and $\mathbf{\Gamma}_b$ have the only meaning of initial orbital elements of particles 1 and 2.

To obtain the equation of evolution of $N_1(h, t)$ let us multiply equation (30) by $d\mathbf{e}d\mathbf{i}$ and integrate. Using equation (16) we can easily do this with the l.h.s. and the first term of the r.h.s. To treat the second term in the r.h.s. of (30) we substitute the distribution function (17) into equation (30) and integrate over $d\mathbf{e}$ using (27) and the following identity:

$$\int \frac{d\mathbf{e}}{4\pi^2\sigma_{e1}^2\sigma_{e2}^2} \exp\left[-\frac{\mathbf{e}_1^2}{2\sigma_{e1}^2} - \frac{\mathbf{e}_2^2}{2\sigma_{e2}^2}\right] = \frac{1}{2\pi(\sigma_{e1}^2 + \sigma_{e2}^2)} \exp\left[-\frac{\mathbf{e}_r^2}{2(\sigma_{e1}^2 + \sigma_{e2}^2)}\right] \equiv \psi_r(\mathbf{e}_r), \quad (31)$$

with $\psi_r(\tilde{\mathbf{e}}_r)$ being the distribution function of *relative* eccentricity as defined by (31). An analogous identity defines the distribution function of relative inclination $\psi_r(\tilde{\mathbf{i}}_r)$. These identities show that the distribution function for the *relative* motion of planetesimals with r.m.s. eccentricities and inclinations σ_{e1}, σ_{i1} and σ_{e2}, σ_{i2} is also Gaussian, with dispersions of the relative eccentricities and inclinations given by

$$\sigma_{er}^2 = \sigma_{e1}^2 + \sigma_{e2}^2, \quad \sigma_{ir}^2 = \sigma_{i1}^2 + \sigma_{i2}^2. \quad (32)$$

Using this result we are able to write the following equation for the evolution of $N_1(h)$:

$$\begin{aligned} \frac{\partial N_1(h)}{\partial t} = & -2|A| \int_0^\infty dm_2 (\mu_1 + \mu_2)^{2/3} \int_{-\infty}^\infty d\tilde{h} |\tilde{h}| \left\{ N_1(h) N_2[h - (\mu_1 + \mu_2)^{1/3} \tilde{h}] \right. \\ & \left. - (\mu_1 + \mu_2)^{4/3} \int d\tilde{\mathbf{e}}_r d\tilde{\mathbf{i}}_r d(\Delta\tilde{\mathbf{\Gamma}}_r) \psi_r(\mathbf{e}_r, \sigma_{er}) \psi_r(\mathbf{i}_r, \sigma_{ir}) N_1(h_1) N_2(h_2) P_r(\tilde{\mathbf{\Gamma}}_r, \Delta\tilde{\mathbf{\Gamma}}_r) \right\}, \end{aligned} \quad (33)$$

where

$$\sigma_{er}^2(h_1, h_2) = \sigma_{e1}^2(h_1) + \sigma_{e2}^2(h_2), \quad \sigma_{ir}^2(h_1, h_2) = \sigma_{i1}^2(h_1) + \sigma_{i2}^2(h_2), \quad (34)$$

$$\begin{aligned} h_1 &= h + D(\Delta\tilde{h}), \quad h_2 = h + D(\Delta\tilde{h}) - (\mu_1 + \mu_2)^{1/3} \tilde{h}, \\ \text{with} \quad D(\Delta\tilde{h}) &= -\frac{\mu_2}{(\mu_1 + \mu_2)^{2/3}} \Delta\tilde{h}. \end{aligned} \quad (35)$$

In the last term of (33) one can perform the integration over $d(\Delta\tilde{\mathbf{\Gamma}}_r)$ using equation (29) [which reduces to simple replacement of $\Delta\tilde{h}$ by $\Delta\tilde{h}_{sc}(\tilde{h}, \tilde{\mathbf{e}}_r, \tilde{\mathbf{i}}_r)$, see equation (29)]. Also it is more natural to use the *reduced* velocity distribution function $\tilde{\psi}_r(\tilde{\mathbf{e}}_r, \tilde{\mathbf{i}}_r)$ defined as

$$\tilde{\psi}_r(\tilde{\mathbf{e}}_r, \tilde{\mathbf{i}}_r) = (\mu_1 + \mu_2)^{4/3} \psi_r(\mathbf{e}_r) \psi_r(\mathbf{i}_r) = \frac{1}{4\pi^2 \tilde{\sigma}_{er}^2 \tilde{\sigma}_{ir}^2} \exp \left[-\frac{\tilde{\mathbf{e}}_r^2}{2\tilde{\sigma}_{er}^2} - \frac{\tilde{\mathbf{i}}_r^2}{2\tilde{\sigma}_{ir}^2} \right], \quad (36)$$

where

$$\tilde{\sigma}_{e,i} = \frac{\sigma_{e,i}}{(\mu_1 + \mu_2)^{1/3}}. \quad (37)$$

Using all these results and definitions one can finally transform equation (33) into

$$\begin{aligned} \frac{\partial N_1(h)}{\partial t} = & -2|A| \int_0^\infty dm_2 (\mu_1 + \mu_2)^{2/3} \int_{-\infty}^\infty d\tilde{h} |\tilde{h}| \left\{ N_1(h) N_2[h - (\mu_1 + \mu_2)^{1/3} \tilde{h}] \right. \\ & \left. - \int d\tilde{\mathbf{e}}_r d\tilde{\mathbf{i}}_r \tilde{\psi}_r(\tilde{\mathbf{e}}_r, \tilde{\mathbf{i}}_r) N_1[h + D(\Delta\tilde{h}_{sc})] N_2[h + D(\Delta\tilde{h}_{sc}) - (\mu_1 + \mu_2)^{1/3} \tilde{h}] \right\}. \end{aligned} \quad (38)$$

The functions $N_{1,2}(h)$ and $\tilde{\psi}_r(\tilde{\mathbf{e}}_r, \tilde{\mathbf{i}}_r)$ are also functions of time. This equation is analogous to the evolution equation (6) of Paper I but now it can also take into account the dependence of scattering properties on the random motion of interacting bodies.

2.5. Evolution of random velocities.

Let us multiply equation (30) by $\mathbf{e}^2 d\mathbf{e} d\mathbf{i}$ and, again, integrate over the whole vector eccentricity and inclination space. Since integrating the l.h.s. of (30) and the first term in its r.h.s. is again trivial [by definition $\int \mathbf{e}^2 f_j(\mathbf{e}, \mathbf{i}) d\mathbf{e} d\mathbf{i} = 2N_j \sigma_e^2$], we concentrate on the second term in the r.h.s. We notice using (31) that the integration of $\psi_1 \psi_2 \mathbf{e}^2 d\mathbf{e}$ reduces to

$$\begin{aligned} & \exp \left[-\frac{\mathbf{e}_r^2}{2(\sigma_{e1}^2 + \sigma_{e2}^2)} \right] \int \frac{\mathbf{e}^2 d\mathbf{e}}{4\pi^2 \sigma_{e1}^2 \sigma_{e2}^2} \exp \left[-\frac{\sigma_{e1}^2 + \sigma_{e2}^2}{2\sigma_{e1}^2 \sigma_{e2}^2} \left(\mathbf{e} - \frac{\mu_2}{\mu_1 + \mu_2} \Delta \mathbf{e}_r - \frac{\sigma_{e1}^2 \mathbf{e}_r}{\sigma_{e1}^2 + \sigma_{e2}^2} \right)^2 \right] \\ = & \psi_r(\mathbf{e}_r) \left[2 \frac{\sigma_{e1}^2 \sigma_{e2}^2}{\sigma_{e1}^2 + \sigma_{e2}^2} + \frac{\sigma_{e1}^4 \mathbf{e}_r^2}{(\sigma_{e1}^2 + \sigma_{e2}^2)^2} + \left(\frac{\mu_2}{\mu_1 + \mu_2} \right)^2 (\Delta \mathbf{e}_r)^2 + 2 \frac{\mu_2}{\mu_1 + \mu_2} \frac{\sigma_{e1}^2}{\sigma_{e1}^2 + \sigma_{e2}^2} \mathbf{e}_r \cdot \Delta \mathbf{e}_r \right] \end{aligned} \quad (39)$$

[Integration over $d\mathbf{i}$ yields $\psi_r(\mathbf{i}_r)$, see equations (31)]. Using again the PlDF in the form (29) we find that integration of the second part of the r.h.s. of (30) gives

$$\begin{aligned} & \int d\tilde{\mathbf{e}}_r d\tilde{\mathbf{i}}_r \tilde{\psi}_r(\tilde{\mathbf{e}}_r, \tilde{\mathbf{i}}_r) N_1[h + D(\Delta\tilde{h}_{sc})] N_2[h + D(\Delta\tilde{h}_{sc}) - (\mu_1 + \mu_2)^{1/3} \tilde{h}] \\ & \times \left[2 \frac{\sigma_{e1}^2 \sigma_{e2}^2}{\sigma_{e1}^2 + \sigma_{e2}^2} + \frac{\sigma_{e1}^4 \mathbf{e}_r^2}{(\sigma_{e1}^2 + \sigma_{e2}^2)^2} + \left(\frac{\mu_2}{\mu_1 + \mu_2} \right)^2 (\Delta \mathbf{e}_{sc})^2 + 2 \frac{\mu_2}{\mu_1 + \mu_2} \frac{\sigma_{e1}^2}{\sigma_{e1}^2 + \sigma_{e2}^2} \mathbf{e}_r \cdot \Delta \mathbf{e}_{sc} \right], \end{aligned} \quad (40)$$

where we have again used function $\tilde{\psi}_r(\tilde{\mathbf{e}}_r, \tilde{\mathbf{i}}_r)$ defined in equation (36), and

$$\Delta \mathbf{e}_{sc}(\tilde{h}, \tilde{\mathbf{e}}_r, \tilde{\mathbf{i}}_r) = (\mu_1 + \mu_2)^{1/3} \Delta \tilde{\mathbf{e}}_{sc}(\tilde{h}, \tilde{\mathbf{e}}_r, \tilde{\mathbf{i}}_r) \quad (41)$$

comes from equation (29).

Using these results we can rewrite the evolution equation of the r.m.s. eccentricity of planetesimals of mass m_1 in the following form:

$$\begin{aligned} \frac{\partial}{\partial t} [2N_1(h)\sigma_{e1}^2(h)] &= -2|A| \int_0^\infty dm_2 (\mu_1 + \mu_2)^{2/3} \int_{-\infty}^\infty d\tilde{h} |\tilde{h}| \\ &\times \left\{ 2\sigma_{e1}^2(h)N_1(h)N_2[h - (\mu_1 + \mu_2)^{1/3}\tilde{h}] - \int d\tilde{\mathbf{e}}_r d\tilde{\mathbf{i}}_r \tilde{\psi}_r(\tilde{\mathbf{e}}_r, \tilde{\mathbf{i}}_r) N_1(h_1)N_2(h_2) \right. \\ &\times \left[2\frac{\sigma_{e1}^2\sigma_{e2}^2}{\sigma_{e1}^2 + \sigma_{e2}^2} + \frac{\sigma_{e1}^4 \mathbf{e}_r^2}{(\sigma_{e1}^2 + \sigma_{e2}^2)^2} + \left(\frac{\mu_2}{\mu_1 + \mu_2}\right)^2 (\Delta \mathbf{e}_{sc})^2 + 2\frac{\mu_2}{\mu_1 + \mu_2} \frac{\sigma_{e1}^2}{\sigma_{e1}^2 + \sigma_{e2}^2} \mathbf{e}_r \cdot \Delta \mathbf{e}_{sc} \right] \Big\} \quad (42) \end{aligned}$$

As usual, it is implied in the last term of this equation that

$$\begin{aligned} \sigma_{e1}^2 &= \sigma_{e1}^2(h_1), \quad \sigma_{e2}^2 = \sigma_{e1}^2(h_2), \quad \text{where now} \\ h_1 &= h + D(\Delta \tilde{h}_{sc}), \quad h_2 = h + D(\Delta \tilde{h}_{sc}) - (\mu_1 + \mu_2)^{1/3}\tilde{h}, \end{aligned} \quad (43)$$

i.e. only the kinematic characteristics of planetesimals of different masses at their locations *before* the encounter are important.

It should be remembered that the integral over $d\tilde{\mathbf{e}}_r d\tilde{\mathbf{i}}_r$ in the r.h.s. of equation (42) cannot be easily evaluated in general, because $\Delta \tilde{h}_{sc}$ and $\Delta \mathbf{e}_{sc}$ depend on both $\tilde{\mathbf{e}}_r$ and $\tilde{\mathbf{i}}_r$. However, if the disk kinematic properties and surface density are homogeneous (i.e. σ_{ej} , σ_{ij} , and N_j are independent of h), then the integration over $d\tilde{\mathbf{e}}_r d\tilde{\mathbf{i}}_r$ can be easily performed on the first two terms in the square brackets of equation (42) and they vanish. Planetesimal eccentricities would then evolve only because of the presence of the last two terms in square brackets in (42). This leads us to the conclusion that these nonvanishing terms [third and fourth in the r.h.s. of (42)] are responsible for the planetesimal *self-stirring* — kinematic heating in the course of gravitational scattering of planetesimals (which is present even in a completely homogeneous disk). The equation of eccentricity evolution in homogeneous disk then assumes the form [using (42)]

$$\frac{\partial \sigma_{e1}^2}{\partial t} = |A| \int_0^\infty dm_2 (\mu_1 + \mu_2)^{2/3} N_2 \left[\left(\frac{\mu_2}{\mu_1 + \mu_2}\right)^2 H_1 + 2\frac{\mu_2}{\mu_1 + \mu_2} \frac{\sigma_{e1}^2}{\sigma_{e1}^2 + \sigma_{e2}^2} H_2 \right] \quad (44)$$

where self-stirring coefficients are defined as

$$H_1 = \int_{-\infty}^\infty d\tilde{h} |\tilde{h}| \int d\tilde{\mathbf{e}}_r d\tilde{\mathbf{i}}_r \tilde{\psi}_r(\tilde{\mathbf{e}}_r, \tilde{\mathbf{i}}_r) (\Delta \mathbf{e}_{sc})^2, \quad H_2 = \int_{-\infty}^\infty d\tilde{h} |\tilde{h}| \int d\tilde{\mathbf{e}}_r d\tilde{\mathbf{i}}_r \tilde{\psi}_r(\tilde{\mathbf{e}}_r, \tilde{\mathbf{i}}_r) (\mathbf{e}_r \cdot \Delta \mathbf{e}_{sc}). \quad (45)$$

Equation (44) and definitions (45) are analogous to equations of evolution of planetesimal eccentricity dispersion derived by other authors for the case of a homogeneous planetesimal disk (Ida 1990; Wetherill & Stewart 1993; SI).

The other terms in the r.h.s. of equation (42), which disappear in the homogeneous disk, must describe the *transport* of horizontal energy due to gravitational scattering, and are caused by (1) the gradients of r.m.s. eccentricity and inclination of planetesimals, and (2) the gradients of planetesimal number density. To the best of our knowledge, this part of eccentricity evolution has never been taken explicitly into account before. In the general case of an inhomogeneous disk (e.g. in the presence of a massive embryo which can naturally induce gradients of disk properties by its gravity) both contributions (self-heating and transport) are important.

In analogous fashion we can write the equation for the inclination evolution:

$$\begin{aligned} \frac{\partial}{\partial t} [2N_1(h)\sigma_{i1}^2(h)] = & -2|A| \int_0^\infty dm_2 (\mu_1 + \mu_2)^{2/3} \int_{-\infty}^\infty d\tilde{h} |\tilde{h}| \\ & \left\{ 2\sigma_{i1}^2(h)N_1(h)N_2[h - (\mu_1 + \mu_2)^{1/3}\tilde{h}] - \int d\tilde{\mathbf{e}}_r d\tilde{\mathbf{i}}_r \tilde{\psi}_r(\tilde{\mathbf{e}}_r, \tilde{\mathbf{i}}_r) N_1(h_1)N_2(h_2) \right. \\ & \times \left[2\frac{\sigma_{i1}^2\sigma_{i2}^2}{\sigma_{i1}^2 + \sigma_{i2}^2} + \frac{\sigma_{i1}^4\mathbf{i}_r^2}{(\sigma_{i1}^2 + \sigma_{i2}^2)^2} + \left(\frac{\mu_2}{\mu_1 + \mu_2}\right)^2 (\Delta\mathbf{i}_{sc})^2 + 2\frac{\mu_2}{\mu_1 + \mu_2} \frac{\sigma_{i1}^2}{\sigma_{i1}^2 + \sigma_{i2}^2} \mathbf{i}_r \cdot \Delta\mathbf{i}_{sc} \right] \Big\} \quad (46) \end{aligned}$$

[the definition of $\Delta\mathbf{i}_{sc}$ is analogous to definition of $\Delta\mathbf{e}_{sc}$ in equation (41)]. We cannot simplify these equations further without making additional assumptions about the scattering properties of planetesimals. These additional approximations will be made in the next section.

3. Fokker-Planck expansion.

Significant simplification can be achieved if gravitational scattering is weak, i.e. changes in quantities characterizing the planetesimal state $\mathbf{\Gamma}$ are small compared with the average values of these quantities before the encounter. Then a Fokker-Planck type expansion can be performed (Lifshitz & Pitaevskii 1981; Binney & Tremaine 1987).

This situation is usually the case for planetesimal-planetesimal interactions because the evolution of the disk’s kinematic properties prior to the emergence of embryos would likely lead to a considerable dynamical “heating” of planetesimal population. This heating brings planetesimal-planetesimal interactions into the dispersion-dominated regime when two interacting planetesimals of types 1 and 2 have $\tilde{\sigma}_{er}^2 + \tilde{\sigma}_{ir}^2 \gg 1$. Then encounters occur at high relative velocities and large-angle scattering is rare. For example, a rocky planetesimal with the radius 50 km at 1 AU has a corresponding mass of $\approx 10^{21}$ g and Hill radius $\approx 10^{-4}$ AU. The critical velocity determining the lower boundary of the dispersion-dominated regime for the gravitational interactions between such planetesimals is about 3 m s^{-1} , which is likely to be exceeded by the velocity of planetesimal

epicyclic motion (e.g. see the results of coagulation simulations by Inaba *et al.* 2001). Thus, the assumption that planetesimal-planetesimal scattering occurs in the dispersion-dominated regime is usually justified in protoplanetary disks.

The necessary condition for the Fokker-Planck expansion is that changes of $h, \mathbf{e}_r, \mathbf{i}_r$ are small compared with their typical magnitudes. For example, for the dimensionless impact parameter h this means that $a\Delta h$ should be much smaller than the typical length scale on which the surface density of planetesimals varies. However, this approximation does not require that the relative distance between interacting bodies ah must be small in comparison to this typical scale. This is in contrast to the local approximation that is often used along with the Fokker-Planck approach. Indeed, when the planetesimal disk is dynamically hot, every planetesimal can interact with all others within the reach of its own epicyclic excursion. Since eccentricity is large in this velocity regime the semimajor axis separation of interacting planetesimals $\sim ae$ could be of the order of or bigger than r_H ; then local approximation (which would assume that semimajor axis separation of interacting planetesimals is small) becomes inapplicable. This complication is explicitly taken into account in our further consideration.

3.1. Surface density evolution equation.

To implement the Fokker-Planck treatment of the surface density evolution equation (38), we first expand the second term of the r.h.s. of equation (38) in a Taylor series in $D(\Delta\tilde{h}_{sc})$ up to the second order. Using the notation $\alpha = (\mu_1 + \mu_2)^{1/3}$ and $h - (\mu_1 + \mu_2)^{1/3}\tilde{h} = h - \alpha\tilde{h}$ we obtain that:

$$\begin{aligned} & \int d\tilde{\mathbf{e}}_r d\tilde{\mathbf{i}}_r \tilde{\psi}_r(\tilde{\mathbf{e}}_r, \tilde{\mathbf{i}}_r) N_1[h + D(\Delta\tilde{h}_{sc})] N_2[h + D(\Delta\tilde{h}_{sc}) - (\mu_1 + \mu_2)^{1/3}\tilde{h}] \\ &= N_1(h) N_2(h - \alpha\tilde{h}) - \frac{\mu_2}{(\mu_1 + \mu_2)^{2/3}} \frac{\partial}{\partial h} \left[\langle \Delta\tilde{h} \rangle N_1(h) N_2(h - \alpha\tilde{h}) \right] \\ &+ \frac{1}{2} \frac{\mu_2^2}{(\mu_1 + \mu_2)^{4/3}} \frac{\partial^2}{\partial h^2} \left[\langle (\Delta\tilde{h})^2 \rangle N_1(h) N_2(h - \alpha\tilde{h}) \right]. \end{aligned} \quad (47)$$

In obtaining this expression we have taken into account that the distribution function $\tilde{\psi}_r$ is a function of $\tilde{\sigma}_{er}^2, \tilde{\sigma}_{ir}^2$, and, thus, should also be expanded in $D(\Delta\tilde{h}_{sc})$. In equation (47) the moments of $\Delta\tilde{h}$ are

$$\langle (\Delta\tilde{h})^\beta \rangle = \int d\tilde{\mathbf{e}}_r d\tilde{\mathbf{i}}_r \left[\Delta\tilde{h}_{sc}(\tilde{h}, \tilde{\mathbf{e}}_r, \tilde{\mathbf{i}}_r) \right]^\beta \tilde{\psi}_r(\tilde{\mathbf{e}}_r, \tilde{\mathbf{i}}_r) \quad (48)$$

where $\beta = 1, 2$, and $\langle (\Delta\tilde{h})^\beta \rangle$ is a function of $\tilde{h}, \tilde{\sigma}_{er}, \tilde{\sigma}_{ir}$. Substituting (47) into (38) we obtain

$$\frac{\partial N_1}{\partial t} = -\frac{\partial}{\partial h} (\Upsilon_1^N N_1) + \frac{\partial^2}{\partial h^2} (\Upsilon_2^N N_1) \quad (49)$$

where

$$\Upsilon_1^N(h) = 2|A| \int_0^\infty dm_2 \mu_2 \int_{-\infty}^\infty d\tilde{h} |\tilde{h}| \langle \Delta\tilde{h} \rangle N_2(h - \alpha\tilde{h}), \quad (50)$$

$$\Upsilon_2^N(h) = |A| \int_0^\infty dm_2 \frac{\mu_2^2}{(\mu_1 + \mu_2)^{2/3}} \int_{-\infty}^\infty d\tilde{h} |\tilde{h}| \langle (\Delta\tilde{h})^2 \rangle N_2(h - \alpha\tilde{h}). \quad (51)$$

Thus, instead of the integro-differential equation (38) we have obtained the partial differential equation (49).

The integration over $d\tilde{h}$ in definitions of transport coefficients Υ_1^N and Υ_2^N takes into account the non-zero range of the planetesimal disk over which a given planetesimal can experience encounters with other bodies (nonlocal scattering). In the local approximation, $\langle \Delta\tilde{h} \rangle$ and $\langle (\Delta\tilde{h})^2 \rangle$ would fall off rapidly with increasing $\tilde{h}$; expanding the integrand in the expression for Υ_1^N in $\tilde{h}$ one could easily derive the *local* Fokker-Planck evolution equation. This equation has already been discussed by Petit & Hénon (1987b) and in Paper I.

3.2. Random velocity evolution equation.

Now we perform the Fokker-Planck expansion of equation (42) for the velocity dispersion. Again, we need to concentrate on the expansion of the terms under the $d\tilde{\mathbf{e}}_r d\tilde{\mathbf{i}}_r$ integral with respect to $D(\Delta\tilde{h}_{sc})$.

The Fokker-Planck expansion usually assumes keeping only the terms of the two leading orders: the first one is linear in the expansion parameter, but can suffer strong cancellation effects when averaging over the scattering outcomes is performed. The second term is quadratic but it does not suffer from cancellation during the averaging and in general has magnitude similar to the first one. In our case terms such as $(\Delta\tilde{h}_{sc})^2 \mathbf{e} \cdot \Delta \mathbf{e}_{sc}$ or $\Delta\tilde{h}_{sc} (\Delta \mathbf{e}_{sc})^2$ are third order in the perturbation and should be neglected (we will comment on the order of their smallness in §4). But terms like $\Delta\tilde{h}_{sc} \mathbf{e} \cdot \Delta \mathbf{e}_{sc}$ or $(\Delta\tilde{h}_{sc})^2 \mathbf{e}^2$ should be kept in our expansion.

Performing this procedure as in §3.1, after some cumbersome but straightforward transformations, we can write the Fokker-Planck equation for eccentricity evolution in the following form:

$$\frac{\partial}{\partial t} [2N_1 \sigma_{e1}^2] = \Upsilon_0^e N_1 - \frac{\partial}{\partial h} (\Upsilon_1^e N_1) + \frac{\partial^2}{\partial h^2} (\Upsilon_2^e N_1) \quad (52)$$

where

$$\begin{aligned} \Upsilon_0^e(h) &= 2|A| \int_0^\infty dm_2 (\mu_1 + \mu_2)^{4/3} \int_{-\infty}^\infty d\tilde{h} |\tilde{h}| N_2(h - \alpha\tilde{h}) \\ &\times \left[\left(\frac{\mu_2}{\mu_1 + \mu_2} \right)^2 \langle (\Delta\tilde{\mathbf{e}})^2 \rangle + 2 \frac{\mu_2}{\mu_1 + \mu_2} \frac{\tilde{\sigma}_{e1}^2}{\tilde{\sigma}_{e1}^2 + \tilde{\sigma}_{e2}^2} \langle \tilde{\mathbf{e}} \cdot \Delta\tilde{\mathbf{e}} \rangle \right], \\ \Upsilon_1^e(h) &= 2|A| \int_0^\infty dm_2 \mu_2 (\mu_1 + \mu_2)^{2/3} \int_{-\infty}^\infty d\tilde{h} |\tilde{h}| N_2(h - \alpha\tilde{h}) \end{aligned} \quad (53)$$

$$\times \left[2 \frac{\tilde{\sigma}_{e1}^2 \tilde{\sigma}_{e2}^2}{\tilde{\sigma}_{e1}^2 + \tilde{\sigma}_{e2}^2} \langle \Delta \tilde{h} \rangle + \frac{\tilde{\sigma}_{e1}^4}{(\tilde{\sigma}_{e1}^2 + \tilde{\sigma}_{e2}^2)^2} \langle \tilde{\mathbf{e}}^2 \Delta \tilde{h} \rangle + 2 \frac{\mu_2}{\mu_1 + \mu_2} \frac{\tilde{\sigma}_{e1}^2}{\tilde{\sigma}_{e1}^2 + \tilde{\sigma}_{e2}^2} \langle (\tilde{\mathbf{e}} \cdot \Delta \tilde{\mathbf{e}}) \Delta \tilde{h} \rangle \right], \quad (54)$$

$$\begin{aligned} \Upsilon_2^e(h) &= |A| \int_0^\infty dm_2 \mu_2^2 \int_{-\infty}^\infty d\tilde{h} |\tilde{h}| N_2(h - \alpha \tilde{h}) \\ &\times \left[2 \frac{\tilde{\sigma}_{e1}^2 \tilde{\sigma}_{e2}^2}{\tilde{\sigma}_{e1}^2 + \tilde{\sigma}_{e2}^2} \langle (\Delta \tilde{h})^2 \rangle + \frac{\tilde{\sigma}_{e1}^4}{(\tilde{\sigma}_{e1}^2 + \tilde{\sigma}_{e2}^2)^2} \langle \tilde{\mathbf{e}}^2 (\Delta \tilde{h})^2 \rangle \right]. \end{aligned} \quad (55)$$

The new scattering coefficients used in these equations are

$$\begin{aligned} \langle (\Delta \tilde{\mathbf{e}})^2 \rangle &= \int \tilde{\psi}_r(\tilde{\mathbf{e}}_r, \tilde{\mathbf{i}}_r) (\Delta \tilde{\mathbf{e}}_{sc})^2 d\tilde{\mathbf{e}}_r d\tilde{\mathbf{i}}_r, \quad \langle \tilde{\mathbf{e}} \cdot \Delta \tilde{\mathbf{e}} \rangle = \int \tilde{\psi}_r(\tilde{\mathbf{e}}_r, \tilde{\mathbf{i}}_r) \tilde{\mathbf{e}}_r \cdot \Delta \tilde{\mathbf{e}}_{sc} d\tilde{\mathbf{e}}_r d\tilde{\mathbf{i}}_r, \\ \langle \tilde{\mathbf{e}}^2 \Delta \tilde{h} \rangle &= \int \tilde{\psi}_r(\tilde{\mathbf{e}}_r, \tilde{\mathbf{i}}_r) \tilde{\mathbf{e}}_r^2 \Delta \tilde{h}_{sc} d\tilde{\mathbf{e}}_r d\tilde{\mathbf{i}}_r, \quad \langle \tilde{\mathbf{e}}^2 (\Delta \tilde{h})^2 \rangle = \int \tilde{\psi}_r(\tilde{\mathbf{e}}_r, \tilde{\mathbf{i}}_r) \tilde{\mathbf{e}}_r^2 (\Delta \tilde{h}_{sc})^2 d\tilde{\mathbf{e}}_r d\tilde{\mathbf{i}}_r, \\ \langle (\tilde{\mathbf{e}} \cdot \Delta \tilde{\mathbf{e}}) \Delta \tilde{h} \rangle &= \int \tilde{\psi}_r(\tilde{\mathbf{e}}_r, \tilde{\mathbf{i}}_r) \Delta \tilde{h}_{sc} (\tilde{\mathbf{e}}_r \cdot \Delta \tilde{\mathbf{e}}_{sc}) d\tilde{\mathbf{e}}_r d\tilde{\mathbf{i}}_r, \end{aligned} \quad (56)$$

These coefficients, like the old ones ($\langle \Delta \tilde{h} \rangle$, $\langle (\Delta \tilde{h})^2 \rangle$) are functions of $\tilde{h}$, $\tilde{\sigma}_{e1}^2$, $\tilde{\sigma}_{e2}^2$, $\tilde{\sigma}_{i1}^2$, $\tilde{\sigma}_{i2}^2$. Analytical expressions for these coefficients in the two-body approximation will be derived in §4. An analogous evolution equation can be written for σ_{i1}^2 .

The first term in the r.h.s. of equation (52) is responsible for the gravitational stirring of eccentricity, while the last two terms describe energy transport and disappear in homogeneous planetesimal disks. The coefficients in (49) and (52) are nonlocal quantities and this is reflected in the presence of an integration over $d\tilde{h}$ in their definitions.

Equations (49)-(51) and (52)-(55) describe self-consistently the evolution of the surface density, eccentricity, and inclination at every point of an inhomogeneous planetesimal disk due to planetesimal-planetesimal gravitational encounters in the dispersion-dominated regime. The principal approximation is that we follow moments rather than whole distribution function. To close this system we only need to supplement it with the expressions for various scattering coefficients appearing in evolution equations.

4. Scattering coefficients in the dispersion-dominated regime.

In this section we derive analytical expressions for the scattering coefficients used in equations (49)-(51) and (52)-(55). In so doing we will always assume that interaction between planetesimals is in the dispersion-dominated regime, for the reasons given at the beginning of §3. In the case of embryo-planetesimal scattering it is less clear that the scattering is dispersion-dominated since the embryo mass and Hill radius are much larger; we consider these additional details in Paper II.

In the dispersion-dominated regime most interactions occur between planetesimals separated by $|\tilde{h}| \gg 1$ (or $|h| \gg r_H$). In this case they can be divided into those that can experience *close*




Fig. 1.— Separation of different scattering regimes in the $e - i$ phase space.

encounters, and those which do not approach each other closely. For the planetesimals of the first type most of the change of their orbital elements occurs near the closest approach point. Moreover, this change occurs so rapidly that one can consider scattering to be *instantaneous* and *local*, i.e. to occur at some point and not over an extended part of the trajectory. These observations considerably simplify the analytical treatment of the problem because one only needs to consider a small region near the encounter point. Clearly, only planetesimals with $\tilde{e} \geq |\tilde{h}|$ fall in this category (all the orbital parameters are relative and we drop subscript “r” in this section). It will turn out that planetesimals in this part of the parameter space ($\tilde{e} \geq |\tilde{h}|$ and arbitrary $\tilde{i}$) are the most important contributors to the scattering coefficients. We will call this part of the $\tilde{\mathbf{e}} - \tilde{\mathbf{i}}$ space “Region 1” and devote §4.1 to its study (see Figure 1).

Some planetesimals cannot approach closely — those with $\tilde{e} < |\tilde{h}|$. This part of the phase space can be split into two more regions: one with $\tilde{i} < |\tilde{h}|$, we will call it “Region 2” (see Figure 1), and the other with $\tilde{i} > |\tilde{h}|$ (“Region 3”). In the Region 2 planetesimals accumulate changes of their orbital parameters along a significant portion of their orbits, which stretches approximately by several times $\tilde{h}$ along the y -direction. In a zeroth approximation, the planetesimal disk in this part of phase space can be treated as not possessing random motion at all. Encounters between planetesimals in Region 2 lead to smaller change in the orbital elements than encounters in Region 1 because close encounters cannot take place in Region 2 (Hasegawa & Nakazawa 1990; SI). In

Region 3 typical distances at which interaction occurs are similar to those of Region 2 but relative velocities are higher (they are magnified by the large value of $\tilde{i}$), which makes scattering in Region 3 weaker than either Region 1 or Region 2.

In general all 3 regions will contribute to the scattering coefficients for a given value of $\tilde{h}$. However, their contributions are not of the same order and their relative magnitudes strongly depend on the value of $\tilde{h}$. Assume first that $|\tilde{h}| \leq \tilde{\sigma}_e$. Then the distribution (17) implies that most of the planetesimals belong to Region 1 (where $\tilde{e} \geq |\tilde{h}|$). Since every individual scattering event in Region 1 is also stronger than those happening in Regions 2 and 3 we can conclude that Region 1 strongly dominates the scattering coefficients when $|\tilde{h}| \leq \tilde{\sigma}_e$. However, if $\tilde{\sigma}_e \leq |\tilde{h}|$ the number of planetesimals in Region 1 of the phase space becomes exponentially small, so that distant encounters (Region 2) will dominate the scattering coefficients.

For planetesimal-planetesimal scattering we are mostly interested in quantities *integrated* over some part of planetesimal disk, such as the coefficients Υ_j^N and Υ_j^e in equations (49) and (52). They are rather insensitive to the details of the distribution of the integrands with $\tilde{h}$. Moreover, one can show that the contribution of distant encounters (Regions 2 and 3) to the integrated quantities is still subdominant compared to those from Region 1 (SI). For this reason we will derive only the contribution to the scattering coefficients from Region 1 of the phase space and will completely neglect distant encounters when discussing the planetesimal-planetesimal interactions. The role of distant encounters in the context of the embryo-planetesimal scattering is more important and will be discussed in Paper II.

4.1. Scattering coefficients due to close encounters.

Since our disk evolution equations use $\tilde{h}$, $\tilde{\mathbf{e}}$, and $\tilde{\mathbf{i}}$ as coordinates and since we are interested in dealing with *inhomogeneous* planetesimal disks it is natural to use the Hill formalism (Ida *et al.* 1993; Tanaka & Ida 1996; SI) to calculate scattering coefficients. This is opposed to the local velocity formalism of Hornung *et al.* (1985) which would be rather awkward for our purposes.

The treatment of the close encounters in the dispersion-dominated regime is significantly simplified by the fact that the interaction has a 3-dimensional character — planetesimals can approach the scatterer from all directions. This is in contrast to distant encounters which are intrinsically 2-dimensional. A characteristic feature of 3-dimensional scattering is that if we order planetesimals by their impact parameter, then equal logarithmic intervals in impact parameter contribute equally to the scattering (Binney & Tremaine 1987). In the dispersion-dominated regime most of this interval corresponds to minimum separations less than r_H . In this case close encounters can be treated by using the *two-body* approximation which neglects the influence of the third body (Sun) on the process of instantaneous local scattering.

We will derive analytical expressions for the scattering coefficients (48) and (56) averaged over $d\tilde{e}d\tilde{i}$. Switching from $\tilde{\mathbf{e}}$ and $\tilde{\mathbf{i}}$ back to $\tilde{e}$, $\tilde{i}$, τ , and ω (absolute values of relative eccentricity and

inclination and corresponding constant phases, see §2.1) we find that $d\tilde{e}d\tilde{i} = \tilde{e}d\tilde{e} \tilde{i}d\tilde{i} d\tau d\omega$. The phases $\tau/2\pi$ and $\omega/2\pi$ are distributed uniformly in the interval $(0, 1)$, while the distribution of $\tilde{e}, \tilde{i}$ has a form [equation (17)]

$$\tilde{\psi}(\tilde{e}, \tilde{i})d\tilde{e}d\tilde{i} = \frac{\tilde{e}d\tilde{e} \tilde{i}d\tilde{i}}{\tilde{\sigma}_e^2 \tilde{\sigma}_i^2} \exp \left[-\frac{\tilde{e}^2}{2\tilde{\sigma}_e^2} - \frac{\tilde{i}^2}{2\tilde{\sigma}_i^2} \right]. \quad (57)$$

We will perform the averaging in two steps: first we average over phases τ and ω (these averages will be denoted by $\langle \dots \rangle_{\tau, \omega}$), and then we average over the absolute values of eccentricity and inclination (these final averages will be denoted by simply $\langle \dots \rangle$).

Thus, initially we fix $\tilde{h}$, $\tilde{e}$, and $\tilde{i}$. The idea of the calculation is to assume the flux and velocity of incoming particles near the location of the scatterer (the planetesimal on which we center the reference frame of relative motion) to be represented by their unperturbed values and use them as an input for the two-body scattering problem. These quantities will be given by their values at the scatterer's location $\mathbf{r} = 0$ which means that we only retain zeroth order terms of their expansion near $\mathbf{r} = 0$. Then the components of relative planetesimal approach velocity $\mathbf{v}$ in the r, φ and z directions are given by evaluating the time derivatives of equations (2) at $x = y = z = 0$:

$$v_x = \pm \Omega r_H \sqrt{\tilde{e}^2 - \tilde{h}^2}, \quad v_y = \Omega r_H \frac{\tilde{h}}{2}, \quad v_z = \pm \Omega r_H \tilde{i}. \quad (58)$$

Everywhere in this section velocities v_x, v_y, v_z are assumed to be defined relative to the *local* circular orbit (i.e. $v_y = \dot{y} - 2Ax$, see §2.1).

Next we replace the variables τ and ω by the impact parameter, l , and the angle ϕ of the orbital plane (see Appendix 8.A of Binney & Tremaine 1987²). We will use l and ϕ as an equivalent set of variables and replace averaging over $\tau/2\pi$ and $\omega/2\pi$ by the averaging over l and ϕ . Since we assume that the planetesimal flux is locally homogeneous around the scatterer the distribution of planetesimal trajectories in ϕ is uniform.

Let us set τ_0 and ω_0 to be the values of τ and ω for which the planetesimal orbit passes through the location of the scatterer $\mathbf{r} = 0$. There are 4 such sets of τ and ω [because v_x and v_z have a sign ambiguity in (58)]. It was demonstrated by Ida *et al.* (1993) that the planetesimal trajectories have impact parameters smaller than l only when the phases τ and ω of their orbits lie within 4 ellipsoids near 4 sets of τ_0, ω_0 in $\tau - \omega$ space. The total surface area covered by these ellipsoids is given by

$$A(l) = \frac{8 \pi l^2}{3 r_H^2} \frac{v}{\Omega r_H \tilde{i} |\tilde{h}| \sqrt{\tilde{e}^2 - \tilde{h}^2}}, \quad (59)$$

where

$$v = \Omega r_H \sqrt{\tilde{e}^2 + \tilde{i}^2 - (3/4)\tilde{h}^2} \quad (60)$$

²In the notation of Binney & Tremaine (1987) $l \equiv b$.

is the magnitude of the planetesimal approach velocity [see (58)]. Thus, particles having impact parameters in the range from l to $l + \Delta l$ cover a surface area $(dA/dl)\Delta l$ in $\tau - \omega$ phase space. This allows us to replace the average of some quantity f with respect to $d\tau d\omega/(4\pi^2)$ by an integration with respect to $ldl d\phi$ using the following conversion:

$$\langle f \rangle_{\tau, \omega} = \int_0^{2\pi} \int_0^{2\pi} f(\tau, \omega) \frac{d\omega d\tau}{4\pi^2} \rightarrow \frac{4}{3\pi} \frac{1}{r_H^2} \frac{v}{\Omega r_H} \frac{1}{\tilde{h} |\tilde{e}^2 - \tilde{h}^2|} \int_0^{l_{max}} l dl \int_0^{2\pi} f(l, \phi) \frac{d\phi}{2\pi}. \quad (61)$$

The upper cutoff distance l_{max} will be specified later. To be precise, the transition represented by (61) is legitimate only when $l \ll r_H$. Some planetesimals pass the scatterer at distances $\geq r_H$ but they contribute only weakly to the overall change of the orbital elements (because of the aforementioned dependence on *logarithmic* and not linear intervals in l). For this reason the errors that arise from using (59) at $|\tau - \tau_0| \sim 1$ or $|\omega - \omega_0| \sim 1$ will hardly affect the integrals in (61).

Using equations (2), (4), and (14) one finds that the various combinations of the changes of the relative orbital parameters that we need can be written as

$$\Delta(\tilde{e}^2) = \frac{1}{\Omega^2 r_H^2} [\Delta(v_x^2) + 4\Delta(v_y^2)], \quad \Delta(\tilde{i}^2) = \frac{1}{\Omega^2 r_H^2} \Delta(v_z^2), \quad (62)$$

$$\tilde{\mathbf{e}} \cdot \Delta \tilde{\mathbf{e}} = \frac{v_x \Delta v_x + 4v_y \Delta v_y}{\Omega^2 r_H^2}, \quad \tilde{\mathbf{i}} \cdot \Delta \tilde{\mathbf{i}} = \frac{v_z \Delta v_z}{\Omega^2 r_H^2} \quad (63)$$

(the collision is assumed to be instantaneous here). Obviously, $\Delta(v_i^2) = 2v_i \Delta v_i + (\Delta v_i)^2$. From equations (4) and (14) we can also derive that $\tilde{h} = (2v_y + \Omega x)/(\Omega r_H)$, meaning that

$$\Delta \tilde{h} = 2 \frac{\Delta v_y}{\Omega r_H} \quad \text{and} \quad (\Delta \tilde{h})^2 = 4 \frac{(\Delta v_y)^2}{\Omega^2 r_H^2}. \quad (64)$$

We also have

$$(\tilde{\mathbf{e}} \cdot \Delta \tilde{\mathbf{e}}) \Delta \tilde{h} = 2 \frac{v_x \Delta v_x \Delta v_y + 4v_y (\Delta v_y)^2}{\Omega^3 r_H^3}, \quad (\tilde{\mathbf{i}} \cdot \Delta \tilde{\mathbf{i}}) \Delta \tilde{h} = 2 \frac{v_z \Delta v_z \Delta v_y}{\Omega^3 r_H^3}. \quad (65)$$

To average these expressions over τ, ω we first integrate them over $d\phi/2\pi$ which means that we need to know $\langle v_i \rangle_\phi$ and $\langle v_i v_j \rangle_\phi$ (here $i = x, y, z$ and $\langle \dots \rangle_\phi$ means averaging over $d\phi/2\pi$). Two-body scattering in terms of coordinates l, ϕ was considered in detail elsewhere (Binney & Tremaine 1987) and we will simply use known results for these averages here:

$$\begin{aligned} \langle \Delta v_i \rangle_\phi &= -\Delta v_\parallel \frac{v_i}{v}, \quad \langle (\Delta v_i)^2 \rangle_\phi = (\Delta v_\parallel)^2 \frac{v_i^2}{v^2} + \frac{1}{2} (\Delta v_\perp)^2 \frac{v^2 - v_i^2}{v^2}, \\ \langle \Delta v_i \Delta v_j \rangle_\phi &= \frac{v_i v_j}{v^2} \left[(\Delta v_\parallel)^2 - \frac{(\Delta v_\perp)^2}{2} \right], \quad i \neq j \end{aligned} \quad (66)$$

where $v = (v_x^2 + v_y^2 + v_z^2)^{1/2}$ and

$$\Delta v_\perp = \frac{2lv^3}{G(m_1 + m_2)} \left[1 + \frac{l^2 v^4}{G^2(m_1 + m_2)^2} \right]^{-1} = 2v \frac{lv^2}{\Omega^2 r_H^3} \left[1 + \left(\frac{lv^2}{\Omega^2 r_H^3} \right)^2 \right]^{-1},$$

$$\Delta v_{\parallel} = 2v \left[1 + \frac{l^2 v^4}{G^2 (m_1 + m_2)^2} \right]^{-1} = 2v \left[1 + \left(\frac{lv^2}{\Omega^2 r_H^3} \right)^2 \right]^{-1}. \quad (67)$$

Here, again, l is the impact parameter.

Using these expressions we find that

$$\begin{aligned} \langle \Delta(\tilde{\mathbf{e}}^2) \rangle_{\phi} &= \frac{1}{\Omega^2 r_H^2} \left[\frac{(\Delta v_{\parallel})^2 - 2v\Delta v_{\parallel}}{v^2} (v_x^2 + 4v_y^2) + \frac{(\Delta v_{\perp})^2}{2v^2} (v_y^2 + 5v_z^2 + 4v_x^2) \right], \\ \langle \Delta(\tilde{\mathbf{i}}^2) \rangle_{\phi} &= \frac{1}{\Omega^2 r_H^2} \left[\frac{(\Delta v_{\parallel})^2 - 2v\Delta v_{\parallel}}{v^2} v_z^2 + \frac{(\Delta v_{\perp})^2}{2v^2} (v_y^2 + v_x^2) \right], \\ \langle \tilde{\mathbf{e}} \cdot \Delta \tilde{\mathbf{e}} \rangle_{\phi} &= -\frac{\Delta v_{\parallel}}{v} \frac{v_x^2 + 4v_y^2}{\Omega^2 r_H^2}, \\ \langle \tilde{\mathbf{i}} \cdot \Delta \tilde{\mathbf{i}} \rangle_{\phi} &= -\frac{\Delta v_{\parallel}}{v} \frac{v_z^2}{\Omega^2 r_H^2}, \\ \langle \Delta \tilde{h} \rangle_{\phi} &= -2 \frac{\Delta v_{\parallel}}{v} \frac{v_y}{\Omega r_H}, \\ \langle (\Delta \tilde{h})^2 \rangle_{\phi} &= \frac{4}{\Omega^2 r_H^2} \left[(\Delta v_{\parallel})^2 \frac{v_y^2}{v^2} + \frac{(\Delta v_{\perp})^2}{2} \frac{v_x^2 + v_z^2}{v^2} \right], \\ \langle (\tilde{\mathbf{e}} \cdot \Delta \tilde{\mathbf{e}}) \Delta \tilde{h} \rangle_{\phi} &= \frac{2v_y}{\Omega^3 r_H^3} \left[(\Delta v_{\parallel})^2 \frac{v_x^2 + 4v_y^2}{v^2} + \frac{(\Delta v_{\perp})^2}{2} \frac{3v_x^2 + 4v_z^2}{v^2} \right], \\ \langle (\tilde{\mathbf{i}} \cdot \Delta \tilde{\mathbf{i}}) \Delta \tilde{h} \rangle_{\phi} &= \frac{2v_y}{\Omega^3 r_H^3} \frac{v_z^2}{v^2} \left[(\Delta v_{\parallel})^2 - \frac{(\Delta v_{\perp})^2}{2} \right]. \end{aligned} \quad (68)$$

We integrate these expressions over l in the range from 0 to l_{max} . In doing this we keep only the leading terms in the logarithmic factor which appears when we integrate $\Delta v_{\parallel}$ and $(\Delta v_{\perp})^2$ over l . In this approximation terms proportional to $(\Delta v_{\parallel})^2$ vanish. As a result we find that

$$\langle \Delta(\tilde{\mathbf{e}}^2) \rangle_{\tau, \omega} = C(\tilde{h}, \tilde{e}, \tilde{i}) \left[2\tilde{e}^2 + 5\tilde{i}^2 - (15/4)\tilde{h}^2 \right], \quad (69)$$

$$\langle \mathbf{e} \cdot \Delta \mathbf{e} \rangle_{\tau, \omega} = -C(\tilde{h}, \tilde{e}, \tilde{i}) \tilde{e}^2 \quad (70)$$

$$\langle \Delta(\tilde{\mathbf{i}}^2) \rangle_{\tau, \omega} = C(\tilde{h}, \tilde{e}, \tilde{i}) \left[\tilde{e}^2 - 2\tilde{i}^2 - (3/4)\tilde{h}^2 \right], \quad (71)$$

$$\langle \mathbf{i} \cdot \Delta \mathbf{i} \rangle_{\tau, \omega} = -C(\tilde{h}, \tilde{e}, \tilde{i}) \tilde{i}^2, \quad (72)$$

$$\langle \Delta \tilde{h} \rangle_{\tau, \omega} = -C(\tilde{h}, \tilde{e}, \tilde{i}) \tilde{h}, \quad (73)$$

$$\langle (\Delta \tilde{h})^2 \rangle_{\tau, \omega} = 4C(\tilde{h}, \tilde{e}, \tilde{i}) \left(\tilde{e}^2 + \tilde{i}^2 - \tilde{h}^2 \right), \quad (74)$$

$$\langle (\mathbf{e} \cdot \Delta \mathbf{e}) \Delta \tilde{h} \rangle_{\tau, \omega} = C(\tilde{h}, \tilde{e}, \tilde{i}) \tilde{h} \left(3\tilde{e}^2 + 4\tilde{i}^2 - 3\tilde{h}^2 \right), \quad (75)$$

$$\langle (\mathbf{i} \cdot \Delta \mathbf{i}) \Delta \tilde{h} \rangle_{\tau, \omega} = -C(\tilde{h}, \tilde{e}, \tilde{i}) \tilde{h} \tilde{i}^2, \quad (76)$$

where

$$C(\tilde{h}, \tilde{e}, \tilde{i}) = \frac{8}{3\pi} \frac{\ln \Lambda}{\tilde{i}|\tilde{h}|\sqrt{\tilde{e}^2 - \tilde{h}^2} \left[\tilde{e}^2 + \tilde{i}^2 - (3/4)\tilde{h}^2 \right]^{3/2}} \left[1 + O\left(\frac{1}{\ln \Lambda}\right) \right], \quad (77)$$

and Coulomb factor

$$\Lambda = l_{max} v^2 / G(m_1 + m_2). \quad (78)$$

The upper cutoff l_{max} is determined by the disk dimensions and we set it to be of the order of the disk thickness $\tilde{i}r_H$ (Stewart & Wetherill 1989; SI). Substituting $l_{max} = \tilde{i}r_H$ and (60) into equation (78) we find that

$$\Lambda = \tilde{i}(c_1 \tilde{e}^2 + c_2 \tilde{i}^2) \gg 1, \quad (79)$$

where $c_{1,2}$ are some constants. By introducing these constants we avoid the need of averaging the logarithmic factors over $\tilde{e}$ and $\tilde{i}$; instead we fix them using numerical data [see Ohtsuki *et al.* (2002) for a similar treatment of Coulomb logarithms]. One can see that all the scattering coefficients that we retain in equations (49)-(51) and (52)-(55) are proportional to $\ln \Lambda \gg 1$. The coefficients of the third and higher orders in the Fokker-Planck expansion do not contain this multiplier. Thus our neglect of these higher-order coefficients has a relative accuracy of $O((\ln \Lambda)^{-1})$.

Using equations (69)-(76) one can easily check that

$$\langle \Delta(\tilde{\mathbf{e}}^2) + \Delta(\tilde{\mathbf{i}}^2) - \frac{3}{4}\Delta(\tilde{h}^2) \rangle_{\tau, \omega} = 0, \quad (80)$$

$$\langle (\tilde{\mathbf{e}} \cdot \Delta \tilde{\mathbf{e}}) \Delta \tilde{h} + (\tilde{\mathbf{i}} \cdot \Delta \tilde{\mathbf{i}}) \Delta \tilde{h} - \frac{3}{4}\tilde{h}(\Delta \tilde{h})^2 \rangle_{\tau, \omega} = 0. \quad (81)$$

The first equation is implied by the conservation of Jacobi constant (15). The second equation is a statement of $\langle \Delta J \Delta \tilde{h} \rangle_{\tau, \omega} = 0$ up to the second order in perturbed quantities [or up to the factors $\sim (\ln \Lambda)^{-1}$].

Now we perform the final step in our programme — we average the scattering coefficients over the Gaussian distribution of relative eccentricity and inclination given by (57). To do this we have to perform a series of straightforward but rather lengthy steps which is described in detail in Appendix B. Coulomb logarithm $\ln \Lambda$ is always treated as a constant, which is justified by its weak dependence on $\tilde{e}$ and $\tilde{i}$. Of course one can do better than this (see Ida *et al.* 1993) but for our present purposes such an accuracy is sufficient.

As a result we obtain the following formulae for our scattering coefficients:

$$\langle \Delta(\tilde{\mathbf{e}}^2) \rangle = K(\tilde{h}, \tilde{\sigma}_e, \tilde{\sigma}_i) \left(-\frac{7}{4}U_0^0 + \frac{7}{8}U_0^1 - \frac{3}{8}U_1^1 \right), \quad (82)$$

$$\langle \tilde{\mathbf{e}} \cdot \Delta \tilde{\mathbf{e}} \rangle = -K(\tilde{h}, \tilde{\sigma}_e, \tilde{\sigma}_i) \left(U_0^0 + \frac{1}{8}U_0^1 + \frac{1}{8}U_1^1 \right), \quad (83)$$

$$\langle \Delta(\tilde{\mathbf{i}}^2) \rangle = K(\tilde{h}, \tilde{\sigma}_e, \tilde{\sigma}_i) \left(\frac{1}{4}U_0^0 - \frac{1}{8}U_0^1 + \frac{3}{8}U_1^1 \right), \quad (84)$$

$$\langle \tilde{\mathbf{i}} \cdot \Delta \tilde{\mathbf{i}} \rangle = -K(\tilde{h}, \tilde{\sigma}_e, \tilde{\sigma}_i) \left(\frac{1}{8} U_0^1 - \frac{1}{8} U_1^1 \right), \quad (85)$$

$$\langle \Delta \tilde{h} \rangle = -\frac{K(\tilde{h}, \tilde{\sigma}_e, \tilde{\sigma}_i)}{\tilde{h}} U_0^0, \quad (86)$$

$$\langle (\Delta \tilde{h})^2 \rangle = K(\tilde{h}, \tilde{\sigma}_e, \tilde{\sigma}_i) U_0^1, \quad (87)$$

$$\langle \tilde{\mathbf{e}}^2 \Delta \tilde{h} \rangle = -\tilde{h} K(\tilde{h}, \tilde{\sigma}_e, \tilde{\sigma}_i) \left(U_0^0 + \frac{1}{8} U_0^1 + \frac{1}{8} U_1^1 \right), \quad (88)$$

$$\langle \tilde{\mathbf{i}}^2 \Delta \tilde{h} \rangle = -\tilde{h} K(\tilde{h}, \tilde{\sigma}_e, \tilde{\sigma}_i) \left(\frac{1}{8} U_0^1 - \frac{1}{8} U_1^1 \right), \quad (89)$$

$$\langle \tilde{\mathbf{e}}^2 (\Delta \tilde{h})^2 \rangle = \tilde{h}^2 K(\tilde{h}, \tilde{\sigma}_e, \tilde{\sigma}_i) \left(U_0^1 + \frac{1}{8} U_0^2 + \frac{1}{8} U_1^2 \right), \quad (90)$$

$$\langle \tilde{\mathbf{i}}^2 (\Delta \tilde{h})^2 \rangle = \tilde{h}^2 K(\tilde{h}, \tilde{\sigma}_e, \tilde{\sigma}_i) \left(\frac{1}{8} U_0^2 - \frac{1}{8} U_1^2 \right), \quad (91)$$

$$\langle (\tilde{\mathbf{e}} \cdot \Delta \tilde{\mathbf{e}}) \Delta \tilde{h} \rangle = \tilde{h} K(\tilde{h}, \tilde{\sigma}_e, \tilde{\sigma}_i) \left(\frac{7}{8} U_0^1 - \frac{1}{8} U_1^1 \right), \quad (92)$$

$$\langle (\tilde{\mathbf{i}} \cdot \Delta \tilde{\mathbf{i}}) \Delta \tilde{h} \rangle = -\tilde{h} K(\tilde{h}, \tilde{\sigma}_e, \tilde{\sigma}_i) \left(\frac{1}{8} U_0^1 - \frac{1}{8} U_1^1 \right). \quad (93)$$

In these expressions we use the following notation:

$$K(\tilde{h}, \tilde{\sigma}_e, \tilde{\sigma}_i) = \frac{4}{3} \frac{\ln \Lambda}{\tilde{\sigma}_e^2 \tilde{\sigma}_i^2} e^{-\tilde{h}^2/(2\tilde{\sigma}_e^2)}, \quad (94)$$

$$U_\rho^\eta(\alpha_e, \alpha_i) \equiv \int_0^\infty dt \frac{t^\eta}{(t+1)^{3/2}} e^{-\frac{1}{2}(\alpha_e^2 + \alpha_i^2)t} I_\rho \left[\frac{1}{2} (\alpha_i^2 - \alpha_e^2) t \right], \quad (95)$$

(I_ρ is a modified Bessel function of order ρ) and it is always assumed in equations (82)-(92) that

$$U_\rho^\eta = U_\rho^\eta \left(\frac{\tilde{h}/\tilde{\sigma}_e}{2\sqrt{2}}, \frac{\tilde{h}/\tilde{\sigma}_i}{2\sqrt{2}} \right). \quad (96)$$

Coefficients $\langle (\Delta \tilde{\mathbf{e}})^2 \rangle$ and $\langle (\Delta \tilde{\mathbf{i}})^2 \rangle$ can be trivially computed from $\langle \Delta(\tilde{\mathbf{e}}^2) \rangle$, $\langle \tilde{\mathbf{e}} \cdot \Delta \tilde{\mathbf{e}} \rangle$, $\langle \Delta(\tilde{\mathbf{i}}^2) \rangle$, and $\langle \tilde{\mathbf{i}} \cdot \Delta \tilde{\mathbf{i}} \rangle$. We also use for the factor inside the Coulomb logarithm the following expression: $\Lambda = \tilde{\sigma}_i(c_1 \tilde{\sigma}_e^2 + c_2 \tilde{\sigma}_i^2) \gg 1$.

We perform some checks and comparisons of our scattering coefficients with the results obtained by other authors. First of all, one can check again in a manner analogous to equations (80) and (81) that (82)-(93) preserve the Jacobi constant. Second we have checked that our formulae for $\langle \Delta(\tilde{\mathbf{e}}^2) \rangle_{\tau, \omega}$, $\langle \mathbf{e} \cdot \Delta \mathbf{e} \rangle_{\tau, \omega}$, $\langle \Delta(\tilde{\mathbf{i}}^2) \rangle_{\tau, \omega}$, $\langle \mathbf{i} \cdot \Delta \mathbf{i} \rangle_{\tau, \omega}$ and $\langle \Delta \tilde{h} \rangle_{\tau, \omega}$ agree with SI and Ida *et al.* (2000). Third, in the case of a homogeneous planetesimal disk integrals of $\langle \Delta(\tilde{\mathbf{e}}^2) \rangle$, $\langle \Delta(\tilde{\mathbf{i}}^2) \rangle$, $\langle \tilde{\mathbf{e}} \cdot \Delta \tilde{\mathbf{e}} \rangle$, and $\langle \tilde{\mathbf{i}} \cdot \Delta \tilde{\mathbf{i}} \rangle$ over $(3/2)|\tilde{h}|d\tilde{h}$ should reproduce the averaged viscous stirring and dynamical friction coefficients $\langle P_{VS} \rangle$, $\langle Q_{VS} \rangle$ and $\langle P_{DF} \rangle$, $\langle Q_{DF} \rangle$ of SI. We numerically checked that this is really the case.

This completes our calculation of the contribution of Region 1 of $\tilde{\mathbf{e}} - \tilde{\mathbf{i}}$ space ($\tilde{e} > |\tilde{h}|$, arbitrary $\tilde{i}$) to the scattering coefficients in the dispersion-dominated regime.

5. Discussion and summary.

We have derived a self-consistent set of equations describing the coupled evolution of the surface density and kinematic properties of a planetesimal disk driven by gravitational encounters between planetesimals. The assumption of a dispersion-dominated velocity regime is used throughout the calculation, which is reasonable for planetesimal disks in their late evolutionary stages. Thus this paper serves as a logical extension of Paper I which was devoted to the study of the shear-dominated velocity regime.

The evolution equations (49) and (52) are of advection-diffusion type; the coefficients entering them are nonlocal which is in contrast with previous results (Paper I; Ohtsuki & Tanaka 2002). This is a natural outcome of the scattering in the dispersion-dominated regime since planetesimals can explore large portions of the disk (compared to their Hill radii) in the course of their epicyclic motion.

We have also derived analytical expressions for the scattering coefficients entering the different terms of evolution equations (50), (51), (53)-(55). The analytical treatment was enabled by the use of the two-body scattering approximation which becomes valid in the dispersion-dominated regime. Our expressions are accurate up to fractional errors $\sim (\ln \Lambda)^{-1}$, where $\ln \Lambda \gg 1$ is a Coulomb logarithm. Following the methods developed in SI and Ohtsuki *et al.* (2002) for the case of homogeneous planetesimal disks it might be possible to improve our calculations by taking subdominant higher order terms into account (they contribute typically at the level $\sim 10\% - 20\%$ and are neglected in our present consideration).

Using this system of evolution equations supplemented with the information about the behavior of the scattering coefficients one can self-consistently study the evolution of inhomogeneous planetesimal disks. Arbitrary distribution of masses of interacting planetesimals is allowed for but the evolution of mass spectrum is not considered in the present study. Thus our equations describe the disk evolution on timescales short compared with the timescale of the mass spectrum evolution. This should be a good assumption for studying effects on the disk caused by the gravity of massive protoplanetary embryos (because they can induce changes of the disk properties on rather short timescales). This deficiency can also be easily remedied in the future when the need to study very long-term disk evolution arises. Our approach can naturally incorporate physical mechanisms other than just gravitational scattering, for example gas drag or migration [see Tanaka & Ida (1999)] and we are going to study their effects in the future.

In the following paper (Paper II) we describe the embryo-planetesimal scattering and derive equations governing this process in various velocity regimes. This would allow us to provide a complete description of the disk evolution caused by both the presence of isolated massive bodies and the continuous sea of planetesimals.

I am indebted to my advisor, Scott Tremaine, for his patient guidance and valuable advice

which I received during the work on this paper. Financial support provided by the Charlotte Elizabeth Procter Fellowship and NASA grant NAG5-10456 is gratefully acknowledged.

REFERENCES

- Binney, J. & Tremaine, S. 1987, *Galactic Dynamics*, Princeton University Press, 1987
- Bodenheimer, P. & Pollack, J.B. 1986, *Icarus*, 67, 391
- Boss, A., Wetherill, G. W., & Haghighipour, N. 2002, *Icarus*, 156, 291
- Butler, R.P., Tinney, C.G., Marcy, G.W., Jones, H.R.A., Penny, A.J., & Apps, K. 2001, *ApJ*, 555, 410
- Goldreich, P. & Lynden-Bell, D. 1965, *MNRAS*, 130, 125
- Goldreich, P. & Tremaine, S. 1980, *ApJ*, 241, 425
- Gradshteyn, I.S. & Ryzhik, I.M. Table of integrals, series, and products, New York: Academic Press, 1980.
- Greenzweig, Y. & Lissauer, J. J. 1992, *Icarus*, 100, 440
- Hasegawa, M. & Nakazawa, K. 1990, *A&A*, 227, 619
- Heinrichsen, I., Walker, H.J., Klaas, U., Sylvester, R.J., & Lemke, D. 1999, *MNRAS*, 304, 589
- Hénon, M. & Petit, J.M. 1986, *Celestial Mechanics*, 38, 67
- Hornung, P., Pellat, R., & Barge, P. 1985, *Icarus*, 64, 295
- Ida, S. 1990, *Icarus*, 88, 129
- Ida, S., Bryden, G., Lin, D. N. C., & Tanaka, H. 2000, *ApJ*, 534, 428
- Ida, S., Kokubo, E., & Makino, J. 1993, *MNRAS*, 263, 875
- Ida, S., & Makino, J. 1992, *Icarus*, 96, 107
- Ida, S., & Makino, J. 1993, *Icarus*, 106, 210
- Inaba, S., Tanaka, H., Nakazawa, K., Wetherill, G. W., & Kokubo, E. 2001, *Icarus*, 149, 235
- Kenyon, S.J. & Luu, J.X. 1998, *AJ*, 115, 2136
- Lifshitz, E. M. & Pitaevskii, L. P. *Physical Kinetics*; New York: Pergamon Press, 1981
- Marcy, G.W., Butler, R.P., & Vogt, S.S. 2000, *ApJL*, 536, 43

- Mayor, M. & Queloz, D. 1995, *Nature*, 378, 355
- Mizuno, H. 1980, *Progr. Theor. Phys.*, 96, 266
- Ohtsuki, K. 1999, *Icarus*, 137, 152
- Ohtsuki, K., Stewart, G. R. & Ida, S. 2002, *Icarus*, 155, 436
- Ohtsuki, K. & Tanaka, H. 2002, *Icarus*, in press
- Petit, J.M. & Hénon, M. 1987a, *A&A*, 173, 389
- Petit, J.M. & Hénon, M. 1987b, *A&A*, 188, 198
- Petit, J.M. & Hénon, M. 1988, *A&A*, 199, 343
- Rafikov, R.R. 2001, *AJ*, 122, 2713, Paper I
- Rafikov, R.R. 2002, submitted to *AJ*, Paper II
- Safronov, V.S. 1972, *Evolution of the Protoplanetary Cloud and Formation of the Earth and Planets*, NASA TT-F-677
- Schneider, G. 2001, AAS Meeting 198, #83.01
- Stevenson, D. J. 1982, *Planet. Space Sci.*, 30, 755
- Stewart, G. & Ida, S. 2000, *Icarus*, 143, 28 (SI)
- Tanaka, H., & Ida, S. 1996, *Icarus*, 120, 371
- Tanaka, H., & Ida, S. 1997, *Icarus*, 125, 302
- Tanaka, H., & Ida, S. 1999, *Icarus*, 139, 350
- Vogt, S.S., Marcy, G.W., Butler, R.P., & Apps, K. 2000, *ApJ*, 536, 902
- Wetherill, G. W. & Stewart, G. R. 1989, *Icarus*, 77, 330
- Wetherill, G. W. & Stewart, G. R. 1993, *Icarus*, 106, 190

A. Calculation of the instantaneous surface density.

Suppose that we know the behavior of the surface density of planetesimal guiding centers $N(h)$ in the whole disk as well as the random velocity distribution function $\psi(e, i, h)$ (we will initially carry out calculations for the case of general velocity distribution function). We want to compute the *instantaneous* surface density $N^{inst}(x_0)$ at some point x_0 , where x_0 is scaled by the reference radius a_0 and is thus dimensionless.

To perform the desired conversion let us denote the number of planetesimals per unit $dv_x dv_y dv_z dx_0 dz$ as $g^{inst}(v_x, v_y, v_z, x_0, z)$ and the same number per unit $dedidhd\tau d\omega$ as $g(e, i, h, \tau, \omega)$. The Jacobian of the transformation between these sets of coordinates [see equations (2) and (13)] is

$$J = \frac{a_0^4}{2} \Omega^3 ei \quad (\text{A1})$$

(z and velocities v_x, v_y, v_z are assumed to be dimensional). Clearly,

$$N^{inst}(x_0) = \int g^{inst}(v_x, v_y, v_z, x_0, z) dv_x dv_y dv_z dz = \int \frac{2g(e, i, h, \tau, \omega)}{a_0^4 \Omega^3 ei} dv_x dv_y dv_z dz. \quad (\text{A2})$$

To compute N^{inst} one needs the following expressions obtained using (2) and (13):

$$e^2 = \frac{v_x^2 + 4v_y^2}{(\Omega a_0)^2}, \quad i^2 = \frac{v_z^2 + \Omega^2 z^2}{(\Omega a_0)^2}, \quad h = x_0 + \frac{2v_y}{\Omega a_0}. \quad (\text{A3})$$

Now we specify a distribution function (17) and find that

$$N^{inst}(x_0) = \frac{1}{2\pi^2 a_0^4 \Omega^3} \int N(h) \frac{dv_x dv_y dv_z dz}{\sigma_e^2(h) \sigma_i^2(h)} \exp \left[-\frac{1}{\Omega^2 a_0^2} \left(\frac{v_x^2 + 4v_y^2}{2\sigma_e^2} + \frac{v_z^2 + \Omega^2 z^2}{2\sigma_i^2} \right) \right], \quad (\text{A4})$$

with h given by (A3). We can easily perform the integration over $dv_x dv_z dz$. Switching also from v_y to h we finally obtain that

$$N^{inst}(x_0) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{dh}{\sigma_e(h)} N(h) \exp \left[-\frac{(x_0 - h)^2}{2\sigma_e^2} \right]. \quad (\text{A5})$$

Note that $N^{inst}(x_0) = N(x_0)$ when $\sigma_e \rightarrow 0$. Also, if $N(h) = \text{const}$, then $N^{inst}(x_0) = N$ according to formula (A5).

B. Averaging over $\tilde{e}$, $\tilde{i}$ distribution.

We illustrate the procedure of averaging the scattering coefficients over the distribution function (57), using the coefficient $\langle \tilde{\mathbf{e}} \cdot \Delta \tilde{\mathbf{e}} \rangle$ as an example. Calculation of other scattering coefficients

is analogous. Using equations (70) and (94) we write:

$$\langle \tilde{\mathbf{e}} \cdot \Delta \tilde{\mathbf{e}} \rangle = -\frac{8}{3\pi} \frac{\ln \Lambda}{|\tilde{h}| \tilde{\sigma}_e^2 \tilde{\sigma}_i^2} \int_{\tilde{h}}^{\infty} \int_0^{\infty} \frac{\tilde{e}^3 d\tilde{e} d\tilde{i} \exp[-\tilde{e}^2/(2\tilde{\sigma}_e^2) - \tilde{i}^2/(2\tilde{\sigma}_i^2)]}{\sqrt{\tilde{e}^2 - \tilde{h}^2} [\tilde{e}^2 + \tilde{i}^2 - (3/4)\tilde{h}^2]^{3/2}}. \quad (\text{B1})$$

The lower limit of the integration over $\tilde{e}$ is set to $\tilde{h}$ because only test bodies with eccentricities bigger than this value can experience close encounters with the reference particle. Introducing new variables x and y by

$$x^2 = \frac{4}{\tilde{h}^2}(\tilde{e}^2 - \tilde{h}^2), \quad y^2 = \frac{4}{\tilde{h}^2}\tilde{i}^2, \quad (\text{B2})$$

we can rewrite the double integral in (B1) as

$$\int \int \dots = 2|\tilde{h}| e^{-\tilde{h}^2/(2\tilde{\sigma}_e^2)} \int_0^{\infty} \int_0^{\infty} \frac{1 + (1/4)x^2}{(1 + x^2 + y^2)^{3/2}} e^{-\alpha_e^2 x^2 - \alpha_i^2 y^2} dx dy, \quad (\text{B3})$$

where

$$\alpha_e^2 = \frac{1}{8} \frac{\tilde{h}^2}{\tilde{\sigma}_e^2}, \quad \alpha_i^2 = \frac{1}{8} \frac{\tilde{h}^2}{\tilde{\sigma}_i^2}. \quad (\text{B4})$$

Making another change of variables $x = \sqrt{r} \cos \theta$, $y = \sqrt{r} \sin \theta$ we obtain that

$$\begin{aligned} \int \int \dots &= |\tilde{h}| e^{-\tilde{h}^2/(2\tilde{\sigma}_e^2)} \int_0^{\infty} \frac{dr}{(1+r)^{3/2}} \exp\left[-\frac{r}{2}(\alpha_e^2 + \alpha_i^2)\right] \\ &\times \int_0^{\pi/2} \left(1 + \frac{r}{8} + \frac{r \cos 2\theta}{8}\right) \exp\left[\frac{r \cos 2\theta}{2}(\alpha_i^2 - \alpha_e^2)\right] d\theta. \end{aligned} \quad (\text{B5})$$

Using the identity (Gradshteyn & Ryzhik 1980)

$$\int_0^{\pi} e^{z \cos t} \cos nt \, dt = \pi I_n(z), \quad (\text{B6})$$

where I_n is a modified Bessel function, and introducing definitions (95) and (96) we finally obtain

$$\int \int \dots = \frac{\pi}{2} |\tilde{h}| e^{-\tilde{h}^2/(2\tilde{\sigma}_e^2)} \left(U_0^0 + \frac{1}{8} U_0^1 + \frac{1}{8} U_1^1 \right). \quad (\text{B7})$$

Using this result and (B1) we finally arrive at equation (83).